

dose without talking to your healthcare provider.

- Do not stop taking pregabalin capsules without talking to your healthcare provider. If you stop taking pregabalin capsules suddenly you may have headaches, nausea, diarrhea, trouble sleeping, increased sweating, or you may feel restless. If you have epilepsy and you stop taking pregabalin capsules suddenly, you may increase your risk of seizures. Tell your healthcare provider about how to stop pregabalin capsules slowly.
- If you miss a dose, take it as soon as you remember. If it is almost time for your next dose, skip the missed dose. Take the next dose at your regular time. Do not take two doses at the same time.
- If you take too much pregabalin, call your healthcare provider or poison control center, or go to the nearest emergency room right away.

What should avoid while taking pregabalin capsules?

- Do not drive a car, work with machines, or do other dangerous activities until you know how pregabalin capsules affects you.
- Do not drink alcohol while taking pregabalin capsules. Pregabalin capsules and alcohol can affect each other and increase side effects such as dizziness and drowsiness.

What are the possible side effects of pregabalin capsules?

Pregabalin capsules may cause serious side effects, including:

- What is the most important information I should know about pregabalin capsules?**
- Muscle problems.** Muscle pain, soreness, or weakness. If you have these symptoms, especially if you feel sick and have a fever, tell your healthcare provider right away.
- Problems with your eyesight, including blurry vision.** Call your healthcare provider if you have any changes in your eyesight.
- Weight gain.** If you lose weight, weight gain may affect the management of your diabetes. Weight gain can also be a serious problem for people with heart problems.
- Fainting "high."**

The most common side effects of pregabalin capsules in adults are:

- dizziness
- weight gain
- trouble concentrating
- blurry vision
- sleepiness
- swelling of hands and feet
- dry mouth

Pregabalin capsules caused skin sores in animal studies. Skin sores did not happen in studies in people. If you have diabetes, you should pay attention to your skin while taking pregabalin capsules and tell your healthcare provider about any sores or skin problems.

Tell your healthcare provider about any side effect that bothers you or that does not go away.

These are not all the possible side effects of pregabalin capsules. For more information, call your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store pregabalin capsules?

- Store pregabalin capsules at 20° to 25°C (68° to 77°F) in its original package.
- Safely throw away any pregabalin capsules that are out of date or no longer needed.

Keep pregabalin capsules and all medicines out of the reach of children.

General information about the safe and effective use of pregabalin capsules

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use pregabalin capsules for a condition for which it was not prescribed. Do not give pregabalin capsules to other people, even if they have the same symptoms you have. It may harm them. You can ask your healthcare provider or pharmacist for information about pregabalin capsules that is written for health professionals.

For more information on Letters Labo Limited at 1-855-495-1905.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store pregabalin capsules?

- Store pregabalin capsules at 20° to 25°C (68° to 77°F) in its original package.
- Safely throw away any pregabalin capsules that are out of date or no longer needed.

Keep pregabalin capsules and all medicines out of the reach of children.

General information about the safe and effective use of pregabalin capsules

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use pregabalin capsules for a condition for which it was not prescribed. Do not give pregabalin capsules to other people, even if they have the same symptoms you have. It may harm them. You can ask your healthcare provider or pharmacist for information about pregabalin capsules that is written for health professionals.

What are the ingredients in pregabalin capsules?

Active ingredients: pregabalin

Inactive ingredients: corn starch, mannitol and talc

Capitule shell, gelatin, sodium lauryl sulfate and titanium dioxide. In addition, 75 mg, 100 mg, 200 mg, 225 mg and 300 mg capsules contain iron oxide.

Imprinting ink: black iron oxide, butyl alcohol, dehydrated alcohol, isopropyl alcohol, potassium hydroxide, propylene glycol, shellac and zinc stearate solution.

Pediatric use information is approved for Pfizer's LYRICA (pregabalin Capsules and Oral Solution products). However, due to Pfizer's marketing exclusivity rights, this drug product is not labeled with that pediatric information.

This Medication Guide has been approved by the U.S. Food and Drug Administration.

Medication Guide available at <http://www.camberpharma.com/medication-guides>

Camber Pharmaceuticals, Inc.
Manufactured by:
Camber Pharmaceuticals, Inc.,
Piscataway, NJ 08854
By: **HYTERIC™**
Notis, LLC Limited
Jensville, Hyderabad - 500 055, India

Revised: 03/2019

4.4 Pediatric Use

4.4.1 Studies Conducted with Children: Postnatal Neurologic, Psychiatric, Behavioral, and Developmental Data

Pharmacokinetics and safety in children have been established.

4.4.2 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.3 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.4 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.5 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.6 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.7 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.8 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.9 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.10 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.11 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.12 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.13 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.14 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.15 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.16 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.17 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.18 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.19 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.20 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.21 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.22 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.23 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.24 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.25 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.26 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.27 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.28 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.29 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

4.4.30 Safety and Efficacy Data

Pharmacokinetics and safety in children have been established.

14.2 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial. The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.3 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.4 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.5 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.6 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.7 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.8 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.9 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.10 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.11 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.12 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.13 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.14 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.15 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.16 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.17 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.18 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.19 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.20 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.21 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.22 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.23 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.24 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.25 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.26 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.27 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.28 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.29 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.30 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.31 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.32 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.33 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.34 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.35 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.36 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.37 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.38 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.39 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.40 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.41 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.42 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.43 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.44 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.45 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.46 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.47 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.48 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.49 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.50 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.51 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.52 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.53 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.54 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.55 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.56 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.57 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.58 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.59 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.60 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.61 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.62 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.63 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.64 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.65 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.66 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.67 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.68 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.69 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.70 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.71 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.72 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.73 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.74 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.75 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.76 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.77 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.78 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.

14.79 Management of Hypertension

The efficacy of pregabalin capsules for management of hypertension was established in a 14-week, double-blind, randomized, placebo-controlled trial.