

-DOSAGE FORMS AND STRENGTHS-

---WARNINGS AND PRECAUTIONS-

Known hypersensitivity to substituted benzimidazoles or any component of the formulation. (4)
Patients receiving rilpivirine-containing products. (4, 7)
Refer to the Contraindications section of the prescribing information for amoxicillin and clarithromycin, when administered in combination

 $\underline{\textit{Gastric Malignancy}}. \ \ \text{In adults, symptomatic response does not preclude the presence of gastric malignancy}. \ \ \text{Consider additional follow-up}$

Acute Tubulointerstitial Nephritis: Discontinue treatment and evaluate patients. (5.2)

Clostridium difficile-Associated Diarrhea: PPI therapy may be associated with increased risk. (5.3)

Bone Fracture: Long-term and multiple daily dose PPI therapy may be associated with an increased risk for osteoporosis-related fractures

Severe Cutaneous Adverse Reactions; Discontinue at the first signs or symptoms of severe cutaneous adverse reactions or other signs of hypersensitivity and consider further evaluation. (5.5)

Cutaneous and Systemic Lupus Erythematosus: Mostly cutaneous; new onset or exacerbation of existing disease; discontinue esomeprazole magnesium and refer to specialist for evaluation. (5.6)

Interaction with Clopidogrel: Avoid concomitant use of esomeprazole magnesium. (5.7)

Cyanocobalamin (Vitamin B-12) Deficiency: Daily long-term use (e.g., longer than 3 years) may lead to malabsorption or a deficiency of

Interaction with St. John's Wort or Rifampin: Avoid concomitant use of esomeprazole magnesium. (5.10, 7)
Interactions with Diagnostic Investigations for Neuroendocrine Tumors: Increased chromogranin A (CgA) levels may interfere with

diagnostic investigations for neuroendocrine tumors, temporarily stop esomeprazole magnesium at least 14 days before assessing CgA

Interaction with Methotrexate: Concomitant use with PPIs may elevate and/or prolong serum concentrations of methotrexate and/or its metabolite, possibly leading to toxicity. With high dose methotrexate administration, consider temporary withdrawal of esomeprazole

<u>Fundic Gland Polyps</u>: Risk increases with long-term use, especially beyond one year. Use the shortest duration of therapy. (5.13)

--- ADVERSE REACTIONS-

To report SUSPECTED ADVERSE REACTIONS, contact Hetero Labs Limited at 1-866-495-1995 or FDA at 1-800-FDA-1088 or www.fda.gov/

Hypomagnesemia and Mineral Metabolism: Reported rarely with prolonged treatment with PPIs. (5.9)

Most common adverse reactions (6.1):

• Adults (≥ 18 years) (≥1%) are: headache, diarrhea, nausea, flatulence, abdominal pain, constipation, and dry mouth.

• Pediatrics (1 to 17 years) (≥2%) are: headache, diarrhea, abdominal pain, nausea, and somnolence.

See full prescribing information for a list of clinically important drug interactions. (7)

See 17 for PATIENT COUNSELING INFORMATION and Medication Guide

• Esomeprazole Magnesium Delayed-Release Capsules: 20 mg and 40 mg. (3)

with esomeprazole magnesium delayed-release capsules. (4)

and diagnostic testing, (5.1)

of the hip, wrist or spine, (5.4)

levels. (5.11, 12.2)

7 DRUG INTERACTIONS

8.1 Pregnancy 8.2 Lactation

10 OVERDOSAGE

11 DESCRIPTION

8.4 Pediatric Use 8.5 Geriatric Use 8.6 Hepatic Impairment

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

12.2 Pharmacodynamics

12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

14.4 Pediatric GERD

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14.4 Feblatio Gallo
14.5 Risk Reduction of NSAID-Associated Gastric Ulcer
14.6 H. pylori Eradication in Adult Patients with Duodenal Ulcer Disease
14.7 Pathological Hypersecretory Conditions, Including Zollinger-Ellison Syndrome, in Adults

Administer esomeprazole magnesium delayed-release capsules orally or via a nasogastric tube, as described below

Swallow esome prazole magnesium delayed-release capsules whole; do not chew or crush the capsules

For patients who have difficulty swallowing capsules, esome prazole magnesium delayed-release capsules can be opened, and the contents sprinkled on applesauce. Use with other foods has not been evaluated and is not recommended.

Open the esome prazole magnesium delayed-release capsule and empty the granules into a 60 mL catheter-tipped syringe

Use the mixture immediately after preparation. Do not administer the granules if they have dissolved or disintegrated.

Attach the catheter-tipped syringe to a nasogastric tube and deliver the contents of the syringe through the nasogastric tube into the

Esomeprazole magnesium delayed-release capsules USP, 20 mg are white opaque size '4' hard gelatin capsule imprinted with "H" on cap and 'E2' on body filled with off white to pale yellow pellets.

Esomeprazole magnesium delayed-release capsules USP, 40 mg are white opaque size '3' hard gelatin capsule imprinted with "H" on cap

Esomeprazole magnesium delayed-release capsules are contraindicated in patients with known hypersensitivity to substituted

benzimidazoles or to any component of the formulation. Hypersensitivity reactions may include anaphylaxis, anaphylactic shock, angioedema, bronchospasm, acute tubulointerstitial nephritis, and urticaria [see Warnings and Precautions (5.2), Adverse Reactions (6.2)]. For information about contraindications of amoxicillin and clarithromycin, indicated in combination with esomeprazole magnesium

delayed-release capsules for H. pylori eradication to reduce the risk of duodenal ulcer recurrence, refer to the Contraindications section of

Proton pump inhibitors (PPIs), including esomeprazole magnesium, are contraindicated in patients receiving rilpivirine-containing

In adults, symptomatic response to therapy with esomeprazole magnesium does not preclude the presence of gastric malignancy. Consider

additional follow-up and diagnostic testing in adult patients who have a suboptimal response or an early symptomatic relapse after completing

Acute tubulointerstitial nephritis (TIN) has been observed in patients taking PPIs and may occur at any point during PPI therapy. Patients

may present with varying signs and symptoms from symptomatic hypersensitivity reactions to non-specific symptoms of decreased renal function (e.g., malaise, nausea, anorexia). In reported case series, some patients were diagnosed on biopsy and in the absence of extrarenal manifestations (e.g., fever, rash or arthralgia). Discontinue esomeprazole magnesium and evaluate patients with suspected acute

ished observational studies suggest that PPI therapy like esomeprazole magnesium may be associated with an increased risk

of Clostridium difficile-associated diarrhea, especially in hospitalized patients. This diagnosis should be considered for diarrhea that does not

Clostridium difficile-associated diarrhea (CDAD) has been reported with use of nearly all antibacterial agents. For more information specific to

antibacterial agents (clarithromycin and amoxicillin) indicated for use in combination with esomeprazole magnesium, refer to Warnings and

Several published observational studies suggest that proton pump inhibitor (PPI) therapy may be associated with an increased risk for

osteoporosis-related fractures of the hip, wrist, or spine. The risk of fracture was increased in patients who received high-dose, defined as multiple daily doses, and long-term PPI therapy (a year or longer). Patients should use the lowest dose and shortest duration of PPI therapy appropriate to the condition being treated. Patients at risk for osteoporosis-related fractures should be managed according to established

Severe cutaneous adverse reactions, including Stevens-Johnson syndrome (SJS) and toxic epidermal necrolysis (TEN), drug reaction with eosinophilia and systemic symptoms (DRESS), and acute generalized exanthematous pustulosis (AGEP) have been reported in association

with the use of PPIs [see Adverse Reactions (6.2)]. Discontinue esomeprazole magnesium at the first signs or symptoms of severe cutaneous

Cutaneous lupus erythematosus (CLE) and systemic lupus erythematosus (SLE) have been reported in patients taking PPIs, including

esomeprazole. These events have occurred as both new onset and an exacerbation of existing autoimmune disease. The majority of PPI-induced

The most common form of CLE reported in patients treated with PPIs was subacute CLE (SCLE) and occurred within weeks to years after

Systemic lupus erythematosus (SLE) is less commonly reported than CLE in patients receiving PPIs. PPI associated SLE is usually milder than

adults to the elderly. The majority of patients presented within days to years after initiating treatment primarily in patients ranging from young adults to the elderly. The majority of patients presented with rash; however, arthralgia and cytopenia were also reported.

Avoid administration of PPIs for longer than medically indicated. If signs or symptoms consistent with CLE or SLE are noted in patients receiving

esomeprazole magnesium, discontinue the drug and refer the patient to the appropriate specialist for evaluation. Most patients improve with discontinuation of the PPI alone in 4 to 12 weeks. Serological testing (e.g., ANA) may be positive and elevated serological test results may take

Avoid concomitant use of esomeprazole magnesium with clopidogrel. Clopidogrel is a prodrug. Inhibition of platelet aggregation by clopidogrel is entirely due to an active metabolite. The metabolism of clopidogrel to its active metabolite can be impaired by use with concomitant medications, such as esomeprazole, that inhibit CYP2C19 activity. Concomitant use of clopidogrel with 40 mg esomeprazole reduces the pharmacological

nuous drug therapy in patients ranging from infants to the elderly. Generally, histological findings were observed without organ in

Patients should use the lowest dose and shortest duration of PPI therapy appropriate to the condition being treated.

Add one tablespoon of applesauce to an empty bowl. The applesauce used should not be hot and should be soft enough to be

* Sections or subsections omitted from the full prescribing information are not listed.

Administer the mixture immediately. Do not chew or crush the granules

Replace the plunger and shake the catheter-tipped syringe vigorously for 15 seconds.

After administering the granules, flush the nasogastric tube with additional water.

Hold the catheter-tipped syringe with the tip up and check for any granules remaining in the tip.

13.2 Animal Toxicology and/or Pharmacology

14.1 Healing of EE in Adults
14.2 Maintenance of Healing of EE in Adults

16 HOW SUPPLIED/STORAGE AND HANDLING

Esomeprazole Magnesium Delayed-Release Capsules

swallowed without chewing.

Mix the granules with the applesauce.

Mix the granules with 50 mL of water.

DOSAGE FORMS AND STRENGTHS

products [see Drug Interactions (7)].

5 WARNINGS AND PRECAUTIONS

TIN [see Contraindications (4)].

5.3 Clostridium difficile-Associated Diarrhea

Precautions section of the corresponding prescribing information

treatment guidelines [see Dosage and Administration (2) and Adverse Reactions (6.2)].

adverse reactions or other signs of hypersensitivity and consider further evaluation

improve [see Adverse Reactions (6.2)].

5.5 Severe Cutaneous Adverse Reactions

5.6 Cutaneous and Systemic Lupus Erythematosus

17 PATIENT COUNSELING INFORMATION

12.4 Microbiology 12.5 Pharmacogenomics

14 CLINICAL STUDIES

8 USE IN SPECIFIC POPULATIONS

magnesium, (5.12, 7)

5.8 Cyanocobalamin (Vitamin B-12) Deficiency
Daily treatment with any acid-suppressing medications over a long period of time (e.g., longer than 3 years) may lead to malabsorption of cyanocobalamin (vitamin B-12) caused by hypo- or achlorhydria. Rare reports of cyanocobalamin deficiency occurring with acid-suppressing

therapy have been reported in the literature. This diagnosis should be considered if clinical symptoms consistent with cyanocobalamin

5.9 Hypomagnesemia and Mineral Metabolism Hypomagnesemia, symptomatic and asymptomatic, has been reported rarely in patients treated with PPIs for at least three months, in most

Hypomagnesemia may lead to hypocalcemia and/or hypokalemia and may exacerbate underlying hypocalcemia in at-risk patients. In most patients, treatment of hypomagnesemia required magnesium replacement and discontinuation of the PPI.

For patients expected to be on prolonged treatment or who take PPIs with medications such as digoxin or drugs that may cause hypomagnesemia (e.g., diuretics), health care professionals may consider monitoring magnesium levels prior to initiation of PPI treatment and

Consider monitoring magnesium and calcium levels prior to initiation of esomeprazole magnesium and periodically while on treatment in patients with a preexisting risk of hypocalcemia (e.g., hypoparathyroidism). Supplement with magnesium and/or calcium, as necessary. If hypocalcemia is refractory to treatment, consider discontinuing the PPI.

5.10 Interaction with St. John's Wort or Rifampin
Drugs which induce CYP2C19 or CYP3A4 (such as St. John's Wort or rifampin) can substantially decrease esomeprazole concentrations [see Drug Interactions (7)]. Avoid concomitant use of esomeprazole magnesium with St. John's Wort or rifampin.

erum chromogranin A (CQA) levels increase secondary to drug-induced decreases in gastric acidity. The increased CQA level may cause false positive esults in diagnostic investigations for neuroendocrine tumors. Healthcare providers should temporarily stop esomeprazole treatment at least 14 days

before assessing CgA levels and consider repeating the test if initial CgA levels are high. If serial tests are performed (e.g., for monitoring), the same

Literature suggests that concomitant use of PPIs with methotrexate (primarily at high dose; see methotrexate prescribing information) may elevate and prolong serum levels of methotrexate and/or its metabolite, possibly leading to methotrexate toxicities. In high-dose methotrexate

5.13 Fundic Gland Polyps
PPI use is associated with an increased risk of fundic gland polyps that increases with long-term use, especially beyond one year. Most PPI users who developed fundic gland polyps were asymptomatic and fundic gland polyps were identified incidentally on endoscopy. Use the shortest duration of PPI therapy appropriate to the condition being treated.

6.1 Clinical Trials Experience
Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be

The safety of esomeprazole magnesium delayed-release capsules was evaluated in over 15,000 patients (aged 18 to 84 years) in clinical trials worldwide including over 8,500 patients in the United States and over 6,500 patients in Europe and Canada. Over 2,900 patients were treated in

commercial laboratory should be used for testing, as reference ranges between tests may vary [see Clinical Pharmacology (12.2)].

administration a temporary withdrawal of the PPI may be considered in some patients [see Drug Interactions (7)].

directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

ADVENSE REAL TURNS

(Following serious adverse reactions are described below and elsewhere in labeling:
Acute Tubulointerstitial Nephritis [see Warnings and Precautions (5.2)]

(Clostridium difficile-Associated Diarrhea [see Warnings and Precautions (5.3)]

Bone Fracture [see Warnings and Precautions (5.4)]

Severe Cutaneous Adverse Reactions [see Warnings and Precautions (5.5)]

Cutaneous and Systemic Lupus Erythematosus [see Warnings and Precautions (5.6)]

Cyanocobalamin (Vitamin B-12) Deficiency [see Warnings and Precautions (5.8)] Hypomagnesemia and Mineral Metabolism [see Warnings and Precautions (5.9)] Fundic Gland Polyps [see Warnings and Precautions (5.13)]

Less common adverse reactions with an incidence of less than 1% are listed below by body system:

Hematologic: anemia, anemia hypochromic, cervical lymphadenopathy, epistaxis, leukocytosis, leukopenia, thrombocytopenia;

Musculoskeletal: arthralgia, arthritis aggravated, arthropathy, cramps, fibromyalgia syndrome, hernia, polymyalgia rheumatica;

Metabolic/Nutritional: glycosuria, hyperuricemia, hyponatremia, increased alkaline phosphatase, thirst, vitamin B12 deficiency, weight increase,

Nervous System/Psychiatric: anorexia, apathy, appetite increased, confusion, depression aggravated, dizziness, hypertonia, nervousness

The following potentially clinically significant laboratory changes in clinical trials, irrespective of relationship to esomeprazole magnesium, were reported in 1% or less of patients: increased creatinine, uric acid, total bilirubin, alkaline phosphatase, ALT, AST, hemoglobin, white blood cell count, platelets, serum gastrin, potassium, sodium, thyroxine and thyroid stimulating hormone [see Clinical Pharmacology (12.2)]. Decreases

Two placebo-controlled studies were conducted in 710 adult patients for the treatment of symptomatic GERD. The most common adverse

The clinical trials of H. pylori eradication of to reduce duodenal ulder recurrence, no additional adverse reactions specific to the combination of esomeprazole magnesium, amoxicillin and clarithromycin were observed and were similar to those observed with esomeprazole magnesium,

amoxicillin, or clarithromycin alone. The most frequently reported adverse reactions for patients who received esomeprazole magnesium lelayed-release capsules, amoxicillin and clarithromycin for 10 days were diarrhea (9%), taste perversion (4%), and abdominal pain (4%). To adverse reactions were observed at higher rates with esomeprazole magnesium, amoxicillin and clarithromycin than were observed with

In clinical trials using of esomeprazole magnesium, amoxicillin and clarithromycin, no additional increased laboratory abnormalities particular

For more information on adverse reactions and laboratory changes with amoxicillin or clarithromycin, refer to Adverse Reactions section of the

The safety of esomeprazole magnesium delayed-release capsules was evaluated in 316 pediatric and adolescent patients aged 1 year to 17 years in four clinical trials for the treatment of symptomatic GERD [see Clinical Studies (14.3)]. In 109 pediatric patients aged 1 year to 11 years, the most frequently reported (at least 1%) treatment-related adverse reactions in these patients were diarrhea (3%), headache (2%) and somnolence

(2%). In 149 pediatric patients aged 12 years to 17 years the most frequently reported adverse reactions (at least 2%) were headache (8%).

The following adverse reactions have been identified during post-approval use of esomeprazole. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug

Metabolism and nutritional disorders: hypomagnesemia (may lead to hypocalcemia and/or hypokalemia) [see Warnings and Precautions (5.9)];

Skin and Subcutaneous Tissue: alopecia, erythema multiforme, hyperhidrosis, photosensitivity, Stevens-Johnson syndrome, toxic epidermal necrolysis (some fatal), drug reaction with eosinophilia and systemic symptoms (DRESS), and acute generalized exanthematous pustulosis

Adverse reactions associated with omeprazole may also be expected to occur with esomeprazole. See the full prescribing information for

Tables 3 and 4 include drugs with clinically important drug interactions and interaction with diagnostics when administered concomitantly with

The effect of PPIs on antiretroviral drugs is variable. The clinical importance and the mechanisms behind these

Decreased exposure of some antiretroviral drugs (e.g., rilpivirine, atazanavir, and nelfinavir) when used

concomitantly with esomeprazole may reduce antiviral effect and promote the development of drug resistance [see Clinical Pharmacology (12.3)].

Increased exposure of other antiretroviral drugs (e.g., saquinavir) when used concomitantly with

esomeprazole may increase toxicity [see Clinical Pharmacology (12.3)].

There are other antiretroviral drugs which do not result in clinically relevant interactions with esomeprazole.

Rilpivirine-containing products: Concomitant use with esomeprazole magnesium is contraindicated

Nelfinavir: Avoid concomitant use with esomeprazole magnesium. See prescribing information for nelfinavir.

Saquinavir: See the prescribing information for saquinavir for monitoring of potential saquinavir-related toxicities

Increased INR and prothrombin time in patients receiving PPIs, including esome prazole, and warfarin concomitantly. Increases in INR and prothrombin time may lead to abnormal bleeding and even death.

Monitor INR and prothrombin time and adjust the dose of warfarin, if needed, to maintain the target INR range.

Concomitant use of esomeprazole with methotrexate (primarily at high dose) may elevate and prolong

serum concentrations of methotrexate and/or its metabolite hydroxymethotrexate, possibly leading to

xate toxicities. No formal drug interaction studies of high-dose methotrexate with PPIs have been

Table 3: Clinically Relevant Interactions Affecting Drugs Co-Administered with Esomeprazole and Interaction with Diagnostics

Atazanavir: See prescribing information for atazanavir for dosing information

Other antiretrovirals: See prescribing information for specific antiretroviral drugs

Consult the labeling of concomitantly used drugs to obtain further information about interactions with PPIs.

conducted [see Warnings and Precautions (5.12)].

[see Contraindications (4)].

esophageal varices, gastric ulcer, gastritis, hernia, benign polyps or nodules, Barrett's esophagus, and mucosal discoloration

nypoesthesia, impotence, insomnia, migraine, migraine aggravated, paresthesia, sleep disorder, somnolence, tremor, vertigo, visual field defect;

hot flushes, fatigue, fever, flu-like disorder, generalized edema, leg edema, malaise, pain, rigors;

Respiratory: asthma aggravated, coughing, dyspnea, larynx edema, pharyngitis, rhinitis, sinusitis;

were seen in hemoglobin, white blood cell count, platelets, potassium, sodium, and thyroxine

reactions that were reported were: diarrhea (4%), headache (4%), and abdominal pain (4%).

Combination Treatment with Esomeprazole Magnesium, Amoxicillin and Clarithromycin

Hepatic: bilirubinemia, hepatic function abnormal, SGOT increased, SGPT increased;

Special Senses: otitis media, parosmia, taste loss, taste perversion;

cases after a year of therapy. Serious adverse events include tetany, arrhythmias, and seizures.

5.11 Interactions with Diagnostic Investigations for Neuroendocrine Tumors

periodically [see Adverse Reactions (6.2)].

ADVERSE REACTIONS

or omeprazole.

Endocrine: goiter;

weight decrease;

polyuria;

Visual: conjunctivitis, vision abnormal

esomeprazole magnesium alone.

<u>Pediatrics</u>

1 Year to 17 Years of Age

Eve: blurred vision:

to these drug combinations were observed.

minal pain (3%), diarrhea (2%), and nausea (2%)

ure. These reports are listed below by body system:

Hepatobiliary: hepatic failure, hepatitis with or without jaundice;

Nervous System: hepatic encephalopathy, taste disturbance; Psychiatric: aggression, agitation, depression, hallucination;

Respiratory, Thoracic, and Mediastinal: bronchospasm

Reproductive System and Breast: gynecomastia, erectile dysfunction;

esomeprazole and instructions for preventing or managing them.

Renal and Urinary: interstitial nephritis;

(AGEP), cutaneous lupus erythematosus.

DRUG INTERACTIONS

Clinical Impact:

Warfarin

Intervention:

Methotrexate

Clinical Impact:

Gastrointestinal: pancreatitis; stomatitis; microscopic colitis; fundic gland polyps

Infections and Infestations: GI candidiasis; Clostridium difficile-associated diarrhea

Musculoskeletal and Connective Tissue: muscular weakness, myalgia, bone fracture;

Immune System: anaphylactic reaction/shock; systemic lupus erythematosu

Blood and Lymphatic: agranulocytosis, pancytopenia

up to 12 months compared to short-term treatmen

Cardiovascular: flushing, hypertension, tachycardia;

Revised: 04/2024

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ESOMEPRAZOLE MAGNESIUM DELAYED-RELEASE CAPSULES safely and effectively. See full prescribing information for ESOMEPRAZOLE MAGNESIUM DELAYED-RELEASE CAPSULES.

Initial U.S. Approval: 1989 (omeprazole)

-INDICATIONS AND USAGE-Esomeprazole magnesium delayed-release capsules are a proton pump inhibitor (PPI).

- Esomeprazole magnesium delayed-release capsules are indicated for the:

 Short-term treatment in the healing of erosive esophagitis (EE) in adults and pediatric patients 12 years to 17 years of age. (1.1) Maintenance of healing of EE in adults. (1.2)
- Short-term treatment of heartburn and other symptoms associated GERD in adults and pediatric patients 12 years to 17 years of age. (1.3) Risk reduction of nonsteroidal anti-inflammatory drugs (NSAID)-associated gastric ulcer in adults at risk for developing gastric ulcers due to age (60 years and older) and/or documented history of gastric ulcers. (1.4)

---DOSAGE AND ADMINISTRATION

Helicobacter pylori eradication in adult patients to reduce the risk of duodenal ulcer recurrence in combination with amoxicillin and

· Long-term treatment of pathological hypersecretory conditions, including Zollinger-Ellison syndrome in adults. (1.6)

Recommended Adult (2.1) and Pediatric Dosage (2.2) Population Healing of EE (1 year and older) EE due to Acid-Mediated GERD (1 month to less than 1 year) 20 mg or 40 mg¹ once daily for 4 to 8 weeks; some patients may require an additional 4 to 8 weeks 12 years to 17 years 20 mg or 40 mg¹ once daily for 4 to 8 weeks Maintenance of Healing of EE 20 mg once daily. Controlled studies do not extend beyond 6 months Treatment of Symptomatic GERD 20 mg once daily once daily for 4 weeks some patients may require an additional 4 weeks 12 years to 17 years 20 mg once daily for 4 weeks Risk Reduction of NSAID-Associated Gastric Ulce 20 mg or 40 mg¹ once daily for up to 6 months² H. pylori Eradication to Reduce the Risk of Duodenal Ulcer Recurrence Esomeprazole magnesium delayed-release capsules 40 mg¹ once daily for 10 days Amoxicillin 1000 mg twice daily for 10 days Clarithromycin 500 mg twice daily for 10 days Pathological Hypersecretory Conditions Including Zollinger-Ellison Syndrome

Starting dosage is 40 mg twice daily4 (varies with the individual patient) as long as clinically indicated ¹ A maximum dosage of 20 mg once daily is recommended for patients with severe liver impairment (Child-Pugh Class C). ² Controlled studies do not extend beyond 6 months.

Refer to the amoxicillin and clarithromycin prescribing information for dosage adjustments in elderly and renally-impaired patients.
 A starting dosage of 20 mg twice daily is recommended for patients with severe liver impairment (Child-Pugh Class C).
 Preparation and Administration Information

Swallow capsules whole; do not crush or chew. For patients who cannot swallow intact capsule, the capsule can be opened, and the contents mixed with applesauce. (2.3)

Opened capsules can be administered through a nasogastric tube. (2.3)

FULL PRESCRIBING INFORMATION: CONTENTS*

- 1 INDICATIONS AND USAGE
- Healing of Erosive Esophagitis (EE) Maintenance of Healing of EE
- Treatment of Symptomatic GERD
 Risk Reduction of Nonsteroidal Anti-Inflammatory Drugs (NSAID)-Associated Gastric Ulcer
 Helicobacter pylori Eradication to Reduce the Risk of Duodenal Ulcer Recurrence
- 1.6 Pathological Hypersecretory Conditions Including Zollinger-Ellison Syndrome
- 2 DOSAGE AND ADMINISTRATION 2.1 Recommended Dosage in Adults by Indication
- Recommended Dosage in Pediatric Patients by Indication Preparation and Administration Instructions
- 3 DOSAGE FORMS AND STRENGTHS 4 CONTRAINDICATIONS
- 5 WARNINGS AND PRECAUTIONS
- Presence of Gastric Malignancy Acute Tubulointerstitial Nephritis Clostridium difficile-Associated Diarrhea
- Bone Fracture
- 5.5 Severe Cutaneous Adverse Reactions
- 5.6 Cutaneous and Systemic Lupus Erythematosus
 5.7 Interaction with Clopidogrel 5.8 Cyanocobalamin (Vitamin B-12) Deficiency
- 5.9 Hypomagnesemia and Mineral Metabolism
 5.10 Interaction with St. John's Wort or Rifampin 5.11 Interactions with Diagnostic Investigations for Neuroendocrine Tumors
- 5.12 Interaction with Methotrexate 5.13 Fundic Gland Polyps
- 6 ADVERSE REACTIONS 6.1 Clinical Trials Experience 6.2 Postmarketing Experience

FULL PRESCRIBING INFORMATION

1.1 Healing of Erosive Esophagitis (EE)

Esomeprazole magnesium delayed-release capsules are indicated for the short-term treatment (4 to 8 weeks) in the healing and symptomatic resolution of diagnostically confirmed EE in adults. For those patients who have not healed after 4 to 8 weeks of treatment, an additional 4- to 8- week course of esomeprazole magnesium delayed-release capsules may be considered.

um delayed-release capsules are indicated for the short-term treatment (4 to 8 weeks) for the healing of EE in pediatric patients 12 years to 17 years of age. 1.2 Maintenance of Healing of EE

Esomeprazole magnesium delayed-release capsules are indicated for the maintenance of healing of EE in adults. Controlled studies do not extend beyond 6 months.

Esomeprazole magnesium delayed-release capsules are indicated for short-term treatment (4 to 8 weeks) of heartburn and other symptoms associated with GERD in adults

Pediatric Patients 12 Years to 17 Years of Age Esomeprazole magnesium delayed-release capsules are indicated for short-term treatment (4 weeks) of heartburn and other symptoms associated with GERD in pediatric patients 12 years to 17 years of age.

1.4 Risk Reduction of Nonsteroidal Anti-Inflammatory Drugs (NSAID)-Associated Gastric Ulcer

Esomeprazole magnesium delayed-release capsules are indicated for the reduction in the occurrence of gastric ulcers associated with continuous NSAID therapy in adult patients at risk for developing gastric ulcers. Patients are considered to be at risk due to their age (60 years and older) and/or documented history of gastric ulcers. Controlled studies do not extend beyond 6 months.

1.5 Helicobacter pylori Eradication to Reduce the Risk of Duodenal Ulcer Recurrence Eradication of *H. pylori* has been shown to reduce the risk of duodenal ulcer recurrence

Esomeprazole magnesium delayed-release capsules in combination with amoxicillin and clarithromycin is indicated for the treatment of adult patients with H. pylori infection and duodenal ulcer disease (active or history of within the past 5 years) to eradicate H. pylori.

In patients who fail therapy, susceptibility testing should be done. If resistance to clarithromycin is demonstrated or susceptibility testing is not possible, alternative antimicrobial therapy should be instituted [see Clinical Pharmacology (12.4) and the prescribing information for clarithromycin].

1.6 Pathological Hypersecretory Conditions Including Zollinger-Ellison Syndrome Esomeprazole magnesium delayed-release capsules are indicated for the long-term treatment of pathological hypersecretory conditions, including Zollinger-Ellison Syndrome, in adults.

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage in Adults by Indication Table 1 shows the recommended adult dosage of esomeprazole magnesium delayed-release capsules by indication.

The duration of esome prazole magnesium delayed-release capsules treatment should be based on available safety and efficacy data specific to the defined indication and dosing frequency and individual patient medical needs. Esomeprazole magnesium delayed-release capsules should

Table 1: Recommended Dosage of Esomeprazole Magnesium Delayed-Release Capsules in Adults by Indicatio

Adult Indication	Recommended Dosage of Esomeprazole Magnesium Delayed- Release Capsules	Treatment Duration		
Healing of EE	20 mg or 40 mg ¹ once daily	4 to 8 weeks ²		
Maintenance of Healing of EE	20 mg once daily	Controlled studies do not extend beyond 6 months		
Treatment of Symptomatic GERD	20 mg once daily	4 weeks; if symptoms do not resolve completely, consider an additional 4 weeks		
Risk Reduction of NSAID-Associated Gastric Ulcer	20 mg or 40 mg ¹ once daily	Controlled studies do not extend beyond 6 months		
H. pylori Eradication to Reduce the Risk of	Esomeprazole magnesium delayed- release capsules 40 mg once daily ¹	10 days		
Duodenal Ulcer Recurrence (Triple Therapy)	Amoxicillin 1000 mg twice daily ³	10 days		
	Clarithromycin 500 mg twice daily ³	10 days		
Pathological Hypersecretory Conditions Including Zollinger-Ellison Syndrome	Starting dosage is 40 mg twice daily ⁴ ; individualize the regimen to patient needs. Dosages of up to 240 mg/day have been administered [see Clinical Studies (14.7)].	As long as clinically indicated		

^{1.} A maximum dosage of 20 mg once daily is recommended for patients with severe liver impairment (Child-Pugh Class C) [see Use in Specific Populations (8.6) Most patients are healed within 4 to 8 weeks. For patients who do not heal after 4 to 8 weeks, an additional 4 to 8 weeks of treatment may be required to achieve healing *[see Clinical Studies (14.1)].* • Refer to the amoxicillin and clarithromycin prescribing information for dosage adjustments in elderly and renally-impaired patients 4 A starting dosage of 20 mg twice daily is recommended for patients with severe liver impairment (Child-Pugh Class C) [see Use in Specific

Populations (8.6)]. 2.2 Recommended Dosage in Pediatric Patients by Indication

Table 2 shows the recommended dosage of esomeprazole magnesium delayed-release capsules in pediatric patients by indication Table 2: Recommended Dosage of Esomeprazole Magnesium Delayed-Release Capsules in Pediatric Patients by Indication

Indication	Patient Age	Recommended Dosage	Duration		
Healing of EE	12 years to 17 years	Esomeprazole magnesium delayed-release capsules: 20 mg or 40 mg once daily	4 to 8 Weeks		
Treatment of Symptomatic GERD	12 years to 17 years	Esomeprazole magnesium delayed-release capsules: 20 mg once daily	4 weeks		
2.3 Preparation and Administration Instructions					

Take esomeprazole magnesium delayed-release capsules at least one hour before meals [see Clinical Pharmacology (12.3)]

d-release could still

Customer/Country | Camber / USA

Dimensions

Antacids may be used concomitantly with esomeprazole magnesium delayed-release capsu Take a missed dose as soon as possible. If it is almost time for the next dose, skip the missed dose and take the next dose at the regular

can side s?"

ease the sleas

See Sium

activity of clopidogrel. When using esomeprazole magnesium consider alternative anti-platelet therapy [see Drug Interactions (7)]

sym ive 4-8 ules

longer to resolve than clinical manifestations.

5.7 Interaction with Clopidogrel

ng d infection the antibi

i of G

ins to cont

ΈΕ

rilpivi treat <u>N</u> levels, blood.

to oreas baby ules plar into feed

Take

you all, y Wort ate).

E B B of as

Use mixim

Effects on maternal bone were observed in pregnant and lactating rats in a pre- and postnatal toxicity study when esomeprazole magnesium was administered at oral doses of 14 to 280 mg/kg/day (about 3.4 to 68 times an oral human dose of 40 mg on a body surface area basis). When rats were dosed from gestational day 7 through weaning on postnatal day 21, a statistically significant decrease in maternal femur weight of up to 14% (as compared to placebo treatment) was observed at doses equal to or greater than 138 mg/kg/day (about 34 times an oral human dose was performed with esomeprazole magnesium at oral doses of 280 mg/kg/day (about 68 times an oral human dose of 40 mg on a body surface area basis) where esomeprazole administration was from either gestational day 7 or gestational day 16 until parturition. When maternal administration was confined to gestation only, there were no effects on bone physeal morphology in the offspring at any age.

mothers not exposed to any proton pump inhibitor during the first trimester. A retrospective cohort study reported on 689 pregnant women exposed to either H2-blockers or omeprazole in the first trimester (134 exposed to omeprazole) and 1,572 pregnant women unexposed to either during the first trimester. The overall malformation rate in offspring born to mothers with first trimester exposure to omeprazole, an H2-blocker, or were unexposed was 3.6%, 5.5%, and 4.1% respectively. Several studies have reported no apparent adverse short-term effects on the infant when single dose oral or intravenous omeprazole was administered to over 200 pregnant women as premedication for cesarean section under general anesthesia productive studies conducted with omeprazole in rats at oral doses up to 138 mg/kg/day (about 34 times an oral human dose of 40 mg on

A pre- and postnatal developmental toxicity study in rats with additional endpoints to evaluate hone development was performed with

of 40 mg on a body surface area basis).

someprazole is the S-isomer of omeprazole and limited data suggest that omeprazole may be present in human milk. There are no clinical data on the effects of esomeprazole on the breastfed infant or on milk production. The developmental and health benefits of breastfeding should be considered along with the mother's clinical need for esomeprazole magnesium delayed-release capsules and any potential adverse effects on the breastfed infant from esomeprazole magnesium delayed-release capsules and any potential adverse effects on the breastfed infant from esomeprazole magnesium delayed-release capsules and any potential adverse effects on the

See Contraindications, Warnings and Precautions in prescribing information for clarithromycin. See Drug Interactions in prescribing information for amoxicillin. patients receiving MMF [see Clinical Pharmacology (12.3)].
See the prescribing information for other drugs dependent on gastric pH for absorption Potentially increased exposure of tacrolimus, especially in transplant patients who are intermediate or poo metabolizers of CYP2C19. Monitor tacrolimus whole blood concentrations and consider reducing the dose, if needed, to maintain therapeutic drug concentrations. See prescribing information for tacrolimus.

Voriconazole Clinical Impact: Increased exposure of esomeprazole [see Clinical Pharmacology (12.3)]. Dose adjustment of esomeprazole magnesium is not normally required. However, in patients with Zollinger-Ellison

Intervention:

<u>Omeprazole</u>

There are no adequate and well-controlled studies with esomeprazole in pregnant women. Esomeprazole is the S-isomer of omeprazole Available epidemiologic data fail to demonstrate an increased risk of major congenital malformations or other adverse pregnancy outcomes with first trimester omeprazole use (see Data). Reproduction studies in rats and rabbits resulted in dose-dependent embryo-lethality at omeprazole doses that were approximately 3.4 to 34 times an oral human dose of 40 mg (based on a body surface area for a 60 kg person).

Teratogenicity was not observed in animal reproduction studies with administration of oral esomeprazole magnesium in rats and rabbits with

Clinical Impact: Decreased exposure of esomeprazole when used concomitantly with strong inducers [see Clinical Pharmacology (12.3)].

St. John's Wort, rifampin: Avoid concomitant use with [see Warnings and Precautions (5.10)].

Ritonavir-containing products: see prescribing information for specific drugs

defects and miscarriage in clinically recognized pregnancies is 2% to 4% and 15% to 20%, respectively. Esomeprazole is the S-isomer of omeprazole. Four epidemiological studies compared the frequency of congenital abnormalities among infants

A population-based retrospective cohort study covering all live births in Denmark from 1996 to 2009, reported on 1,800 live births whose mothers used omeprazole during the first trimester of pregnancy and 837,317 live births whose mothers did not use any proton pump inhibitor. The overall rate of birth defects in infants born to mothers with first trimester exposure to omeprazole was 2.9% and 2.6% in infants born to

A small prospective observational cohort study followed 113 women exposed to omeprazole during pregnancy (89% with first trimeste exposures). The reported rate of major congenital malformations was 4% in the omeprazole group, 2% in controls exposed to non-teratogens, and 2.8% in disease paired controls. Rates of spontaneous and elective abortions, preterm deliveries, gestational age at delivery, and mean birth

a body surface area basis) and in rabbits at doses up to 69.1 mg/kg/day (about 34 times an oral human dose of 40 mg on a body surface area a basis) during organogenesis did not disclose any evidence for a teratogenic potential of omergrazole. In rabbits, omegrazole in a dose range of 6.9 to 69.1 mg/kg/day (about 3.4 to 34 times an oral human dose of 40 mg on a body surface area basis) administered during organogenesis produced dose-related increases in embryo-lethality, fetal resorptions, and pregnancy disruptions. In rats, dose-related embryo/fetal toxicity and mg/kg/day (about 68 times an oral human dose of 40 mg on a body surface area basis) or in rabbits at oral doses up to 86 mg/kg/day (about 41 times an oral human dose of 40 mg on a body surface area basis) administered during organogenesis

rats treated with oral doses of esomeprazole magnesium at doses equal to or greater than 138 mg/kg/day (about 34 times an oral human dos of 40 mg on a body surface area basis).

Clinical Impact: Pharmacology (12.3)]. tervention Consider reducing the dose of cilostazol to 50 mg twice daily. See prescribing information for cilostazol. Digoxin Clinical Impact: Potential for increased exposure of digoxin [see Clinical Pharmacology (12.3)] Monitor digoxin concentrations and adjust the dose, if needed, to maintain therapeutic drug concentrations. Se prescribing information for digoxin. Combination Therapy with Clarithromycin and Amoxicillin Concomitant administration of clarithromycin with other drugs can lead to serious adverse reactions, including Clinical Impact: potentially fatal arrhythmias, and are contraindicated. tervention. Drugs Dependent on Gastric pH for Absorption (e.g., iron salts, erlotinib, dasatinib, nilotinib, mycophenolate mofetil, ketoconazole/ traconazole) Clinical Impact: Esomeprazole can reduce the absorption of other drugs due to its effect on reducing intragastric acidity Mycophenolate mofetil (MMF): Co-administration of omeprazole, of which esomeprazole is an enantiomer, in healthy subjects and in transplant patients receiving MMF has been reported to reduce the exposure to the active metabolite, mycophenolic acid (MPA), possibly due to a decrease in MMF solubility at an increased gastric pH. The clinical relevance of reduced MPA exposure on organ rejection has not been established in transplant patients receiving esomeprazole magnesium and MMF. Use esomeprazole magnesium with caution in transplant ntervention

comparison with the approved dose of clopidogrel.

anti-platelet therapy [see Warnings and Precautions (5.7)].

A temporary withdrawal of esomeprazole magnesium may be considered in some patients receiving high-dose

Concomitant use of esomeprazole 40 mg resulted in reduced plasma concentrations of the active metabolite of clopidogrel and a reduction in platelet inhibition [see Clinical Pharmacology (12.3)].

There are no adequate combination studies of a lower dose of esomeprazole or a higher dose of clopidogrel in

Avoid concomitant use with esomeprazole magnesium delayed-release capsules Consider use of alternative

Increased exposure of citalopram leading to an increased risk of QT prolongation [see Clinical Pharmacology (12.3)].

Increased exposure of cilostazol and one of its active metabolites (3,4-dihydro-cilostazol) [see Clinical

Limit the dose of citalopram to a maximum of 20 mg per day. See prescribing information for citalopram.

Antiretrovirals

Clopidogre

Citalopram

Clinical Impact

ntervention:

Cilostazol

Clinical Impact:

 ${\bf 2C19~Substrates~(e.g.,~clopidogrel,~citalopram,~cilostazol)}\\$

Tacrolimus Clinical Impact: Interactions with Investigations of Neuroendocrine Tumors Serum chromogranin A (CgA) levels increase secondary to PPI-induced decreases in gastric acidity. The increased CgA level may cause false positive results in diagnostic investigations for neuroendocrine tumors [see Warnings and Precautions (5.11), Clinical Pharmacology (12.2)]. Discontinue esomeprazole magnesium at least 14 days before assessing CgA levels and consider repeating the test if initial CgA levels are high. If serial tests are performed (e.g. for monitoring), the same commercial laboratory should be used for testing, as reference ranges between tests may vary.

The safety in the treatment of healing of EE in adults was assessed in four randomized comparative clinical trials, which included 1,240 patients who received esomeprazole magnesium delayed-release capsules 20 mg once daily, 2,434 patients on esomeprazole magnesium delayed-release capsules 40 mg once daily, and 3,008 patients on omeprazole 20 mg once daily. The most frequently occurring adverse reactions (at least 1%) in all three groups were headache (5,5%, 5%, and 3,8%, respectively) and diarrhea (no difference among the three groups). Nausea, flatulence, abdominal pain, constipation, and dry mouth occurred at similar rates among patients taking esomeprazole magnesium Interaction with Secretin Stimulation Test Body as a Whole: abdomen enlarged, allergic reaction, asthenia, back pain, chest pain, substernal chest pain, facial edema, peripheral edema, Hyper-response in gastrin secretion in response to secretin stimulation test, falsely suggesting gastrinoma. Clinical Impact: ntervention: Discontinue esomeprazole magnesium 4 weeks prior to testing [see Clinical Pharmacology (12.2)] False Positive Urine Tests for THC There have been reports of false positive urine screening test for tetrahydrocannabinol (THC) in patients Clinical Impact: Gastrointestinal: bowel irregularity, constipation aggravated, dyspepsia, dysphagia, dysplasia GI, epigastric pain, eructation, esophageal disorder, frequent stools, gastroenteritis, GI hemorrhage, GI symptoms not otherwise specified, hiccup, melena, mouth disorder, pharynx disorder, rectal disorder, serum gastrin increased, tongue disorder, tongue edema, ulcerative stomatitis, vomiting; An alternative confirmatory method should be considered to verify positive results Table 4: Clinically Relevant Interactions Affecting Esomeprazole When Co-Administered with Other Drugs CYP2C19 or CYP3A4 Inducers

syndrome, who may require higher doses, dosage adjustment may be considered. Skin and Appendages: acne, angioedema, dermatitis, pruritus, pruritus ani, rash, rash erythematous, rash maculo-papular, skin inflammation, sweating increased, urticaria; See prescribing information for voriconazole. USE IN SPECIFIC POPULATIONS 8.1 Pregnancy Urogenital: abnormal urine, albuminuria, cystitis, dysuria, fungal infection, hematuria, micturition frequency, moniliasis, genital moniliasis,

doses about 68 times and 42 times, respectively, an oral human dose of 40 mg (based on a body surface area basis for a 60 kg person). Changes in bone morphology were observed in offspring of rats dosed through most of pregnancy and lactation at doses equal to or greater than approximately 34 times an oral human dose of 40 mg. When maternal administration was confined to gestation only, there were no effects Endoscopic findings that were reported as adverse reactions include: duodenitis, esophageil stricture, esophageal ulceration, on bone physeal morphology in the offspring at any age ($see\ Data$). The incidence of adverse reactions during 6-month trials for the maintenance of healing of EE with esomeprazole magnesium delayed-release The estimated background risks of major birth defects and miscarriage for the indicated population are unknown. All pregnancies have a capsules 20 mg once daily was similar to placebo. There were no differences in types of adverse reactions seen during maintenance treatment und risk of birth defect, loss or other adverse outcomes. In the U.S. general pop tion, the estimated background risk of major birt

> born to women who used omeprazole during pregnancy with the frequency of abnormalities among infants of women exposed to H2-receptor A population-based retrospective cohort epidemiological study from the Swedish Medical Birth Registry, covering approximately 99% of pregnancies, from 1995 to 1999, reported on 955 infants (824 exposed during the first trimester with 39 of these exposed beyond first trimester, and 131 exposed after the first trimester) whose mothers used omeprazole during pregnancy. The number of infants exposed in utero to omeprazole that had any malformation, low birth weight, low Apgar score, or hospitalization was similar to the number observed in this population. The number of infants born with ventricular septal defects and the number of stillborn infants was slightly higher in the omeprazoleexposed infants than the expected number in this population.

weight were similar among the groups.

postnatal developmental toxicity were observed in offspring resulting from parents treated with omeprazole at 13.8 to 138 mg/kg/day (about 3.4 to 34 times an oral human dose of 40 mg on a body surface area basis), administered prior to mating through the lactation pe No effects on embryo-fetal development were observed in reproduction studies with esomeprazole magnesium in rats at oral doses up to 280

esomeprazole magnesium at oral doses of 14 to 280 mg/kg/day (about 3.4 to 68 times an oral human dose of 40 mg on a body surface area basis). Neonatal/early postnatal (birth to weaning) survival was decreased at doses equal to or greater than 138 mg/kg/day (about 34 times an oral human dose of 40 mg on a body surface area basis). Body weight and body weight gain were reduced and neurobehavioral or general developmental delays in the immediate post-weaning timeframe were evident at doses equal to or greater than 69 mg/kg/day (about 17 times an oral human dose of 40 mg on a body surface area basis). In addition, decreased femur length, width and thickness of cortical bone, decreased thickness of the tibial growth plate and minimal to mild bone marrow hypocellularity were noted at doses equal to or greater than 14 mg/kg/ day (about 3.4 times an oral human dose of 40 mg on a body surface area basis). Physeal dysplasia in the femur was observed in offspring of

A pre- and postnatal development study in rats with esomeprazole strontium (using equimolar doses compared to esomeprazole magnesium study) produced similar results in dams and pups as described above. A follow up developmental toxicity study in rats with further time points to evaluate pup bone development from postnatal day 2 to adulthood

Risk Summary 8.4 Pediatric Use Healing of EE Pediatric Patients 1 Year to 17 Years of Age The safety and effectiveness of esomeprazole magnesium delayed-release capsules have been established in pediatric patients 12 years to 17 years for short-term treatment (4 to 8 weeks) for healing of EE. Use of esomeprazole magnesium delayed-release capsules for this indication is supported by evidence from adequate and well-controlled studies in adults with additional safety and pharmacokinetic data in pediatric patients 1 year to 17 years of age. The safety profile in pediatric patients 1 year to 17 years of age was similar to adults [see Adverse Reactions (6.1), Clinical Pharmacology (12.3), Clinical Studies (14.4)]. Symptomatic GERD Pediatric Patients 1 Year to 17 Years of Age The safety and effectiveness of esomeprazole magnesium delayed-release capsules have been established in pediatric patients 12 years to 17

Black 00

Note: Pharma Code, Material Code, Product Name and 2D Data Matrix Orientation will be change based on Machine folding feasibility at vendor.

500 x 600 mm (Book Fold: 35 x 35 mm)

(**hip** may

Bible Paper 28 GSM **Pantone Colours** Version No.

years of age for the short-term treatment (4 weeks) of heartburn and other symptoms associated with GERD. Use of esomeprazole magnesium for this indication is supported by evidence from adequate and well-controlled studies in adults with additional safety and pharmacokinetic data in pediatric patients 1 year to 17 years of age. The safety profile in pediatric patients 1 year to 17 years of age was similar to adults [see Adverse Reactions (6.1), Clinical Pharmacology (12.3), Clinical Studies (14.4)].

The safety and effectiveness of esomeprazole magnesium for the treatment of symptomatic GERD in pediatric patients less than 1 year of age have not been established.

Other Conditions
The safety and effectiveness of esome prazole magnesium for the risk reduction of NSAID-associated gastric ulcer, H. pylori eradication to reduce the risk of duodenal ulcer recurrence and treatment of pathological hypersecretory conditions have not been established in pediatric patients.

Juvenile Animal Toxicity Studies
In a juvenile rat toxicity study, esomeprazole was administered with both magnesium and strontium salts at oral doses about 34 to 68 times a daily human dose of 40 mg based on body surface area. Increases in death were seen at the high dose, and at all doses of esomeprazole, there were decreases in body weight, body weight gain, femur weight and femur length, and decreases in overall growth [see Nonclinical Toxicology (13.2)].

8.5 Geriatric UseOf the total number of patients who received esomeprazole magnesium in clinical trials, 1459 were 65 to 74 years of age and 354 patients were 75 years of age and older.

No overall differences in safety and efficacy were observed between the elderly and younger individuals, and other reported clinical experience has not identified differences in responses between elderly and younger patients, but greater sensitivity of some older individuals cannot be ruled out.

In patients with severe hepatic impairment (Child-Pugh Class C) exposure to esomeprazole substantially increased compared to healthy subjects. Dosage modification of esomeprazole magnesium delayed-release capsules is recommended for patients with severe hepatic impairment for the healing of EE, risk reduction of NSAID-associated gastric ulcer, *H. pylori* eradication to reduce the risk of duodenal ulcer recurrence, and pathological hypersecretory conditions including Zollinger-Ellison Syndrome [see Dosage and Administration (2.1), Clinical Pharmacology (12.3)]. In patients with mild to moderate liver impairment (Child-Pugh Classes A and B), no dosage adjustment is necessary.

10 OVERDOSAGE Manifestations in patients exposed to omeprazole, the racemic mixture, at doses up to 2,400 mg (120 times the usual recommended clinical dose) include confusion, drowsiness, blurred vision, tachycardia, nausea, diaphoresis, flushing, headache, dry mouth, and other adverse reactions similar to those seen at recommended dosages. See the full prescribing information for omeprazole for complete safety information. No specific antidote for esomeprazole is known. Since esomeprazole is extensively protein bound, it is not expected to be removed by dialysis

In the event of overdosage, treatment should be symptomatic and supportive. If over-exposure occurs, call your Poison Control Center at 1-800-222-1222 for current information on the management of poisoning or overdosage and the poison of the management of poisoning or overdosage and the poison of the management of poisoning or overdosage and the poison of the management of poison of the poison of the poison of the management of poison of the poison o

The active ingredient in the esomeprazole magnesium delayed-release capsules, USP for oral administration is 1H-Benzimidazole, 5-methoxy-2-[(S)-[(4-methoxy-3,5-dimethyl-2-pyridinyl]methyl] Sulfinyl], Magnesium Salt (2:1) Trihydrate. Esomeprazole is S-enantiomer of omeprazole. (Initial U.S. approval of esomeprazole magnesium: 2001). Its molecular formula is C₃₄H₃₆MgN₆O₆S_{2.3}H₂O with molecular weight of 767.17 as a trihydrate. The structural formula is:

The magnesium salt is a white to slightly colored powder. It contains 3 moles of water. Slightly soluble in methanol, insoluble in water and in n-Heptane. The stability of esomeprazole magnesium is a function of pH; it rapidly degrades in acidic media, but it has acceptable stability under

Esomeprazole magnesium is supplied in delayed-release capsules. Each delayed-release capsule contains 20 mg, or 40 mg of esomeprazole (equivalent to 22.25 mg esomeprazole magnesium trihydrate, USP) or 40 mg of esomeprazole (equivalent to 44.5 mg esomeprazole magnesium trihydrate, USP) in the form of enteric-coated granules with the following inactive ingredients: glyceryl monostearate, hydroxy propyl cellulose, hypromellose, magnesium stearate, methacrylic acid ethyl acrylate copolymer, polysorbate 80, simethicone, sugar spheres (contains sucrose and starch), talc and triethyl citrate. The capsule shells have the following inactive ingredients: gelatin, titanium dioxide and sodium lauryl sulfate The printing ink contains shellac, propylene glycol, strong ammonia solution, black iron oxide and potassium hydroxide

12 CLINICAL PHARMACOLOGY

Esomeprazole belongs to a class of antisecretory compounds, the substituted benzimidazoles, that suppress gastric acid secretion by specific inhibition of the H-r/K-ATPase enzyme system at the secretory surface of the gastric partial cell. Esomeprazole is protonated and converted in the acidic compartment of the parietal cell forming the active inhibitor, the achiral sulphenamide. Because this enzyme system is regarded as the acid (proton) pump within the gastric mucosa, esomeprazole has been characterized as a gastric acid-pump inhibitor, in that it blocks the final step of acid production. This effect is dose-related and leads to inhibition of both basal and stimulated acid secretion irrespective of the stimulus

The effect of esomeprazole on intragastric pH was determined in adult patients with symptomatic GERD in two separate studies. In the first study of 36 patients, esomeprazole magnesium 40 mg and 20 mg delayed-release capsules were administered once daily over 5 days as shown

Table 5: Effect of Esomeprazole on Intragastric pH on Day 5 (N=36) Following Once Daily Dosing of Esomeprazole Magnesium Delayed-Release Capsules in Adult Patients with Symptomatic GERD

Parameter	Esomeprazole Magnesium Delayed-Release Capsules			
	40 mg once daily	20 mg once daily		
% Time Gastric pH >41 (Hours)	70%² (16.8 h)	53% (12.7 h)		
Coefficient of variation	26%	37%		
Median 24 Hour pH	4.92	4.1		
Coefficient of variation	16%	27%		

² p< 0.01 esomeprazole magnesium 40 mg vs. Esomeprazole magnesium 20 mg

In a second study, the effect on intragastric pH of esomeprazole magnesium 40 mg delayed-release capsules administered once daily over a five-day period was similar to the first study, (% time with pH > 4 was 68% or 16.3 hours).

The effect of esomeprazole on serum gastrin concentrations was evaluated in approximately 2,700 patients in clinical trials of oral esomeprazole for up to 8 weeks and in over 1,300 patients for up to 12 months. The mean fasting gastrin level increased in a dose-related manner. The increase in serum gastrin concentrations reached a plateau within two to three months of therapy and returned to baseline levels within four weeks after

discontinuation of therapy. may cause false positive results in diagnostic investigations for neuroendocrine tumors [see Warnings and Precautions (5.11)]

Human gastric biposy specimens have been obtained from more than 3,000 patients (both pediatrics and adults) treated with omeprazole in long-term clinical trials. The incidence of ECL cell hyperplasia in these studies increased with time; however, no case of ECL cell carcinoids,

dysplasia, or neoplasia has been found in these patients [see Nonclinical Toxicology (13.1)]

In over 1,000 patients treated with oral esomeprazole (10 mg, 20 mg or 40 mg/day) for up to 12 months, the prevalence of ECL cell hyperplasia increased with time and dose. No patient developed ECL cell carcinoids, dysplasia, or neoplasia in the gastric mucosa. Endocrine Effects

Esomeprazole had no effect on thyroid function in adults when given esomeprazole magnesium 20 mg or 40 mg delayed-release capsules once daily for 4 weeks. Other effects of esomeprazole on the endocrine system were assessed in studies of omeprazole. Oral doses of omeprazole 30 mg or 40 mg once daily for 2 to 4 weeks had no effect on carbohydrate metabolism, circulating levels of parathyroid hormone, cortisol, estradiol,

Esomeprazole magnesium delayed-release capsules and esomeprazole magnesium delayed-release oral suspension, showed similar bioavailability after a single dose (40 mg) administration in 94 healthy male and female subjects under fasting conditions. After oral administration, peak plasma levels (Cmix) of esomeprazole occur at approximately 1.5 hours (Tmix). The Cmix increases proportionally when the dose is increased, and there is a three-fold increase in the area under the plasma concentration-time curve (AUC) from 20 to 40 mg. At repeated once-daily dosing with 40 mg, the systemic bioavailability is approximately 90% compared to 64% after a single dose of 40 mg. The mean exposure (AUC) to esomeprazole increases from 4.32 micromol*hr/L on Day 1 to 11.2 micromol*hr/L on Day 5 after 40 mg once daily dosing.

food intake compared to fasting conditions [see Dosage and Administration (2.3)]. The pharmacokinetics of esomeprazole in adult patients with symptomatic GERD following repeated once daily administration of 20 mg and 40 mg esomeprazole magnesium delayed-release capsules over Table 6: Geometric Mean (95% CI) Pharmacokinetic Parameters of Esomeprazole on Day 5 Following Once Daily Dosing of Esomeprazole

The AUC after administration of a single 40 mg dose of esomeprazole magnesium delayed-release capsules is decreased by 43% to 53% after

lagnesium Delayed-Release Capsules in Adult Patients with Symptomatic GERD					
	Esomeprazole Magnesium D	Esomeprazole Magnesium Delayed-Release Capsules			
Parameter ¹ (CV)	40 mg once daily (n=36)	20 mg once daily (n=36)			
AUC (micromol·h/L)	12.6 (42%)	4.2 (59%)			
C _{max} (micromol/L)	4.7 (37%)	2.1 (45%)			
T _{max} (hours)	1.6	16			

Values represent the geometric mean, except the T_{max} , which is the arithmetic mean; CV = Coefficient of variatio

Esomeorazole is a time-dependent inhibitor of CYP2C19, resulting in autoinhibition and nonlinear pharmacokinetics. The systemic exposure increases in a more than dose proportional manner after multiple oral doses of esomeprazoles. Compared to the first dose, the systemic exposure (C_{max} and AUC_{0.249}) at steady state following once a day dosing increased by 43% and 90%, respectively, compared to after the first dose for the 20 mg dose and increased by 95% and 159%, respectively, for the 40 mg dose.

Distribution

Esomeprazole is 97% bound to plasma proteins. Plasma protein binding is constant over the concentration range of 2 to 20 micromol/L. The apparent volume of distribution at steady state in healthy subjects is approximately 16 L.

Esomeprazole is extensively metabolized in the liver by the cytochrome P450 (CYP) enzyme system. The metabolites of esomeprazole lack antisecretory activity. The major part of esomeprazole's metabolism is dependent upon the CYP2C19 isoenzyme, which forms the hydroxy and desmethyl metabolites. The remaining amount is dependent on CYP3A4 which forms the sulphone metabolite.

The plasma elimination half-life of esomeprazole is approximately 1 to 1.5 hours. Less than 1% of parent drug is excreted in the urine. Approximately 80% of an oral dose of esomeprazole is excreted as inactive metabolites in the urine, and the remainder is found as inactive metabolites in the feces.

Combination Therapy with Amoxicillin and Clarithromycin
Esomeprazole magnesium delayed-release capsules 40 mg once daily was given in combination with amoxicillin 1000 mg twice daily for a day to 17 healthy male and female subjects. The mean steady state AUC and C_{max} of esomeprazole increased by 70% and 18%, respectively during combination therapy compared to treatment with esomeprazole magnesium delayed-release capsules alone. The observed increase in esomeprazole exposure during co-administration with amoxicillin and clarithromycin is not expected to be clinically relevant.

The pharmacokinetic parameters for amoxicillin and clarithromycin were similar during combination therapy and administration of each drug alone. However, the mean AUC and Cmax for 14-hydroxyclarithromycin increased by 19% and 22%, respectively, during combination therap compared to treatment with clarithromycin alone. This increase in exposure to 14-hydroxyclarithromycin is not considered to be clinically relevan

The AUC and C_{max} values of esomeprazole were slightly higher (25% and 18%, respectively) in the elderly as compared to younger subjects at steady state. This increase in exposure is not considered clinically relevant.

The pharmacokinetic parameters following repeated dose administration of esomeprazole magnesium 1 mg/kg once daily for 7 to 8 days in 1 month to 11-month-old infants with GERD are summarized in Table 7. Table 7: Summary of Esomeprazole Pharmacokinetic Parameters Following Once Daily Dosing of Oral Esomeprazole Magnesium for 7 to 8

Days in 1 Month to 1 Year Old Infants with GERD

Parameter	Esomeprazole Magnesium 1 mg/kg Orally Once Daily		
AUC (micromol·h/L) (n=7) ¹	3.51		

Oss,max (IIIIOIOIIIOI/L) (II=13)	0.67
t _{1/2} (h) (n=8) ¹	0.93
t _{max} (h) (n=15) ²	3
1. Geometric mean	
² Median	
Subsequent pharmacokinetic simulation analyses	showed that for padiatric nations 1 month to 11 months of age a decade regimen of 2.5 mo

Esomeprazole Magnesium 1 mg/kg Orally Once Daily

Subsequent pharmacokinetic simulation analyses showed that for pediatric patients 1 month to 11 months of age, a dosage regimen of 2.5 mg once daily (body weight 3 to 5 kg), 5 mg once daily (body weight more than 5 to 7.5 kg) and 10 mg once daily for (body weight more than 7.5 to 12 kg) would achieve comparable steady-state plasma exposures (AUC) to that observed with 10 mg once daily in patients 1 year to 11 year of age and 20 mg once daily in patients 12 years to 18 years of age, as well as adults. Apparent clearance (CL/F) increases with age in pediatric patients with GERD from 1 month to 2 years of age

1 Year to 11 Years of Age The pharmacokinetics of esomeprazole were studied in pediatric patients with GERD aged 1 year to 11 years. Following once daily dosing with esomeprazole magnesium delayed-release oral suspension for 5 days, the total exposure (AUC) for the 10 mg dosage in patients aged 6 years to 11 years was similar to that seen with the 20 mg dosage in adults and adolescents aged 12 years to 17 years. The total exposure for the 10 mg dosage in patients aged 1 year to 5 years was approximately 30% higher than the 10 mg dosage in patients aged 6 years to 11 years. The total exposure for the 20 mg dosage in patients aged 6 years to 11 years was higher than that observed with the 20 mg dosage in patients aged 12 years to 17 years and adults, but lower than that observed with the 40 mg dosage in 12 to 17 year-olds and adults. See Table 8. Table 8: Summary of Esomeprazole Pharmacokinetic Parameters Following Once Daily Dosing of Esomeprazole magnesium for Delayed Release Oral Suspension for 5 Days in 1 Year to 11 Year Old Patients with GERD

	Esomeprazole magnesium For Delayed-Release Oral Suspension				
	1 Year to 5 Years	6 Years to	11 Years		
Parameter	10 mg once daily (N = 8)	10 mg once daily (N = 7)	20 mg once daily (N = 6)		
AUC (micromol.h/L) ¹	4.83	3.7	6.28		
C _{max} (micromol/L) ¹	2.98	1.77	3.73		
t _{max} (h) ²	1.44	1.79	1.75		
t ₁₅ λz (h) ¹	0.74	0.88	0.73		
CI/F (L/h) 1	5.99	7.84	9.22		

Geometric mean

Parameter

The pharmacokinetics of esomeprazole magnesium were studied in 28 adolescent patients with GERD aged 12 to 17 years inclusive, in a single center study. Patients were randomized to receive esomeprazole magnesium 20 mg or 40 mg once daily for 8 days. Mean Cmax and AUC values of esomeprazole were not affected by body weight or age; and more than dose-proportional increases in mean Cmax and AUC values were observed between the two dose groups in the study. Overall, esomeprazole magnesium pharmacokinetics in adolescent patients aged 12 to 17 years were similar to those observed in adult patients with symptomatic GERD. See Table 9.

Table 9: Comparison of Esomeprazole Pharmacokinetic Parameters Following Once Daily Dosing of Esomeprazole Magnesium Delayed-Release Capsules in Pediatric Patients 12 Years to 17 Years with GERD and Adults with Symptomatic GERD

Esomeprazole Magnesium Delayed-Release Capsules					
12 Years to 17 Years (N=28)		Adults (N=36)			
20 mg once daily for 8 days	40 mg once daily for 8 days	20 mg once daily 40 mg on for 5 days for 5			
3.65	13.86	4.2	12.6		
1.45	5.13	2.1	4.7		
2	1.75	1.6	1.6		
0.82	1.22	1.2	1.5		
	20 mg once daily for 8 days 3.65 1.45	12 Years to 17 Years (N=28) 20 mg once daily for 8 days 3.65 13.86 1.45 5.13 2 1.75	12 Years to 17 Years (N=28) Adults		

Male and Female Patients

1. Data obtained from two independent studie

The AUC and C_{max} values of esomeprazole were slightly higher (13%) in females than in males at steady state when dosed orally. This increase in exposure is not considered clinically relevant.

The pharmacokinetics of esomeprazole magnesium delayed-release capsules in patients with renal impairment are not expected to be altered relative to healthy subjects as less than 1% of esomeprazole is excreted unchanged in urine.

orally once daily to patients with mild (Child-Pugh Class A, n=4), moderate (Child-Pugh Class B, n=4), and severe (Child-Pugh Class C, n=4)

epatic impairment were compared to those obtained in 36 male and female GERD patients with normal liver function. In patients with mild and moderate hepatic impairment, the AUCs were within the range that could be expected in patients with normal liver function. In patients with severe hepatic impairment the AUCs were 2 to 3 times higher than in the patients with normal liver function [see Use in Specific Populations (8.6)]. <u>Drug Interaction Studies</u> *Effect of Esomeprazole/Omeprazole on Other Drugs*

 $\textit{In vitro} \ \text{and} \ \textit{in vivo} \ \text{studies} \ \text{have shown that esome} \ \text{prazole} \ \text{is not likely to inhibit CYPs 1A2, 2A6, 2C9, 2D6, 2E1} \ \text{and } \ 3A4.$

For some antiretroviral drugs, such as rilpivirine, atazanavir and nelfinavir, decreased serum concentrations have been reported when given together with omeprazole [see Drug Interactions (7)].

Following multiple doses of rilpivirine (150 mg, daily) and omeprazole (20 mg, daily), AUC was decreased by 40%, C_{max} by 40%, and Cmin by 33% for rilpivirine [see Contraindications (4)].

Following multiple doses of nelfinavir (1250 mg, twice daily) and omeprazole (40 mg daily), AUC was decreased by 36% and 92%, Cmax by 37% and one prazole (40 mg daily), AUC was decreased by 36% and 92%, Cmax by 37% and 500 mg daily), AUC was decreased by 36% and 92%, Cmax by 37% and 500 mg daily), AUC was decreased by 36% and 92%, Cmax by 37% and 500 mg daily), AUC was decreased by 36% and 92%, Cmax by 37% and 500 mg daily), AUC was decreased by 36% and 92%, Cmax by 37% and 500 mg daily), AUC was decreased by 36% and 92%, Cmax by 37% and 500 mg daily), AUC was decreased by 36% and 92%, Cmax by 37% and 500 mg daily), AUC was decreased by 36% and 92%, Cmax by 37% and 500 mg daily), AUC was decreased by 36% and 92%, Cmax by 37% and 500 mg daily), AUC was decreased by 36% and 92%, Cmax by 37% and 500 mg daily), AUC was decreased by 36% and 92%, Cmax by 37% and 500 mg daily), AUC was decreased by 36% and 92%, Cmax by 37% and 500 mg daily), AUC was decreased by 36% and 500 mg daily), AUC was decreased by 36% and 500 mg daily), AUC was decreased by 36% and 500 mg daily), AUC was decreased by 36% and 500 mg daily), AUC was decreased by 36% and 500 mg daily), AUC was decreased by 36% and 500 mg daily), AUC was decreased by 36% and 500 mg daily), AUC was decreased by 36% and 500 mg daily). and 89% and Cmin by 39% and 75% respectively for nelfinavir and M8.

Following multiple doses of atazanavir (400 mg, daily) and omeprazole (40 mg, daily, 2 hours before atazanavir), AUC was decreased by 94%, Cmax by 96%, and Cmin by 95%.

Following multiple dosing of saquinavir/ritonavir (1000/100 mg) twice daily for 15 days with omeprazole 40 mg daily co-administered days 11 to 15. The AUC was increased by 82%, C_{max} by 75%, and C_{min} by 106%. The mechanism behind this interaction is not fully elucidated.

Clopidogrel dosage for 28 days) alone and with esomeprazole (40 mg orally once daily at the same time as clopidogrel) for 29 days. Exposure to the active metabolite of clopidogrel was reduced by 35% to 40% over this time period when clopidogrel and esomeprazole were administered together. Pharmacodynamic parameters were also measured and demonstrated that the change in inhibition of platelet aggregation was related to the change in the exposure to clopidogrel active metabolite [see Warnings and Precautions (5.7) and Drug Interactions (7)].

Administration of omeprazole 20 mg twice daily for 4 days and a single 1000 mg dose of MMF approximately one hour after the last dose of omeprazole to 12 healthy subjects in a cross-over study resulted in a 52% reduction in the C_{max} and 23% reduction in the AUC of MPA [see Drug

Omeprazole acts as an inhibitor of CYP2C19. Omeprazole, given in doses of 40 mg daily for one week to 20 healthy subjects in cross-over study, increased Cmax and AUC of cilostazol by 18% and 26% respectively. The Cmax and AUC of one of the active metabolites, 3,4-dihydro-cilostazol, which has 4 to 7 times the activity of cilostazol, were increased by 29% and 69%, respectively. Co-administration of cilostazol with omegrazole s expected to increase concentrations of cilostazol and the above mentioned active metabolite [see Drug Interactions (7)].

Co-administration of esomeprazole 30 mg and diazepam, a CYP2C19 substrate, resulted in a 45% decrease in clearance of diazepam. Increased plasma levels of diazepam were observed 12 hours after dosing and onwards. However, at that time, the plasma levels of diazepam were below the therapeutic interval, and thus this interaction is unlikely to be of clinical relevance.

Concomitant administration of omegrazole 20 mg once daily and digoxin in healthy subjects increased the bioavailability of digoxin by 10% (30%) in two subjects) [see Drug Interactions (7)].

Concomitant administration of esomeprazole and either naproxen (non-selective NSAID) did not identify any clinically relevant changes in the pharmacokinetic profiles of these NSAIDs.

Effect of Other Drugs on Esomeprazole/Omeprazole St. John's Wort

On the August St. St. John's Wort (300 mg three times daily for 14 days) significantly decreased the systemic exposure of omeprazole in CYP2C19 poor metabolizers (Cmax and AUC both decreased by 38%) and extensive metabolizers (Cmax and AUC decreased by 50% and 44%, respectively) [see Drug Interactions (7)].

Concomitant administration of omeprazole and voriconazole (a combined inhibitor of CYP2C19 and CYP3A4) resulted in more than doubling of the omeprazole exposure. When voriconazole (400 mg every 12 hours for one day, followed by 200 mg once daily for 6 days) was given with omeprazole (40 mg once daily for 7 days) to healthy subjects, the steady-state C_{max} and AUC₀₋₂₄ of omeprazole significantly increased: an average of 2 times (90% Cl: 1.8, 2.6) and 4 times (90% Cl: 3.3, 4.4), respectively, as compared to when omeprazole was given without

Co-administration of esome prazole with oral contraceptives, diazepam, phenytoin, quinidine, naproxen (non-selective NSAID) did not seem to change the pharmacokinetic profile of esomeprazole.

Esome prazole magnesium, amoxicillin, and clarithromycin triple therapy has been shown to be active against most strains of Helicobacter pylori (H. pylori) in vitro and in clinical infections [see Indications and Usage (1) and Clinical Studies (14)].

licobacter pvlori: Susceptibility testing of H. pylori isolates was performed for amoxicillin and clarithromycin using agar dilution methodology

and minimum inhibitory concentrations (MICs) were determined. Pretreatment Resistance: Clarithromycin pretreatment resistance rate (MIC \geq 1 mcg/mL) to H. Pylori was 15% (66/445) at baseline in all treatment groups combined. A total of > 99% (394/395) of patients had H. Pylori isolates that were considered to be susceptible (MIC \leq 0.25

mcg/mL) to amoxicillin at baseline. One patient had a baseline $H.\ pylori$ isolate with an amoxicillin MIC = 0.5 mcg/mL. ${\it Clarith romycin Susceptibility Test Results and Clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility results and clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility results and clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility results and clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility results and clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility results and clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility results and clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility results and clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility results and clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility results and clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility results and clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility results and clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility results and clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility results and clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility results and clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility results and clinical/Bacteriologic Outcomes:} \ {\it The baseline H. pylori clarith romycin susceptibility rom$

the H. pylori eradication results at the Day 38 visit are shown in Table 10:

elayed-Release Capsules 40 mg once daily, Amoxicillin 1000 mg twice daily and Clarithromycin 500 mg twice daily for 10 days)						
Clarithromycin	H. pylori negative	H. pylori positive				

Pretreatment	, .	(Eradicated)	Pos	(Not Eradicated) Post-treatment susceptibility results		
			S ²	2	R ²	No MIC
Susceptible ²	182	162	4	0	2	14
Intermediate ²	1	1	0	0	0	0
Resistant ²	29	13	1	0	13	2
		t-treatment clarithromycin susceptibilit e (I) MIC = 0.5 mcg/mL, Resistant (R)				

Patients not eradicated of *H. pylori* following triple therapy with esomeprazole magnesium, amoxicillin and clarithromycin will likely have clarithromycin resistant H. pylori isolates. Therefore, clarithromycin susceptibility testing should be done, when possible. Patients with clarithromycin resistant *H. pylori* should not be re-treated with a clarithromycin-containing regimen.

Amoxicillin Susceptibility Test Results and Clinical/Bacteriological Outcomes: In patients treated with esomeprazole magnesium, amoxicillin and clarithromycin in clinical trials, 83% (176/212) of the patients who had pretreatment amoxicillin susceptible MICs (\leq 0.25 mcg/mL) were eradicated of *H. pylori*, and 17% (36/212) were not eradicated of *H. pylori*. Of the 36 natients who were not eradicated of *H. ovlori*. 16 had no post-treatment susceptibility test results and 20 had post-treatment *H. pylori* isolate with amovicillin susceptible MICs. Fifteen of the patients who were not eradicated of *H. pylori* also had post-treatment *H. pylori* isolates with clarithromycin resistant MICs. There were no patients with *H. pylori* isolates who developed treatment emergent resistance to amovicillin.

Effects on Gastrointestinal Microbial Ecology: Decreased gastric acidity due to any means, including proton pump inhibitors, increases gastric counts of bacteria normally present in the gastrointestinal tract. Treatment with PPIs may lead to slightly increased risk of gastrointestinal infections such as Salmonella and Campylobacter and possibly Clostridium difficile in hospitalized patients

CYP2C19, a polymorphic enzyme, is involved in the metabolism of esome prazole. The CYP2C19*1 allele is fully functional while the CYP2C19*2 and *3 alleles are nonfunctional. There are other alleles associated with no or reduced enzymatic function. Patients carrying two fully functional alleles are extensive metabolizers and those carrying two loss-of-function alleles are poor metabolizers. The systemic exposure to esome prazole raries with a patient's metabolism status: poor metabolizers > intermediate metabolizers > extensive metabolizers. Approximately 3% of Caucasians and 15 to 20% of Asians are CYP2C19 poor metabolizers.

Systemic esomeorazole exposures were modestly higher (approximately 17%) in CYP2C19 intermediate metabolizers (IM: n=6) compared to extensive metabolizers (EM; n=17) of CYP2C19. Similar pharmacokinetic differences were noted across these genotypes in a study of Chinese healthy subjects that included 7 EMs and 11 IMs. There is very limited pharmacokinetic information for poor metabolizers (PM) from these studies. At steady state following once daily administration of esomeprazole 40 mg, the ratio of AUC in poor metabolizers to AUC in the rest of the population (EMs) is approximately 1.5. This change in exposure is not considered clinically meaningful.

13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
The carcinogenic potential of esomeprazole magnesium was assessed using studies of omeprazole, of which esomeprazole is an enanti-In two 24-month oral carcinogenicity studies in rats, omeprazole at daily doses of 1.7, 3.4, 13.8, 44, and 140.8 mg/kg/day (about 0.4 to 34 times the human dose of 40 mg/day expressed on a body surface area basis) produced gastric ECL cell carcinoids in a dose-related manner in both male and female rats, the incidence of this effect was markedly higher in female rats, which had higher blood levels of omeprazole. Gastric carcinoids seldom occur in the untreated rat. In addition, ECL cell hyperplasia was present in all treated groups of both sexes. In one of these studies, female rats were treated with 13.8 mg omeprazole/kg/day (about 3.4 times the human dose of 40 mg/day on a body surface area basis) for 1 year, then followed for an additional year without the drug. No carcinoids were seen in these rats. An increased incidence of treatment-related ECL cell hyperplasia was observed at the end of 1 year (94% treated vs. 10% controls). By the second year the difference between treated and control rats was much smaller (46% vs. 26%) but still showed more hyperplasia in the treated group. Gastric adenocarcinoma was seen in one rat (2%). No similar tumor was seen in male or female rats treated for 2 years. For this strain of rat no similar tumor has been noted listorically, but a finding involving only one tumor is difficult to interpret. A 78-week mouse carcinogenicity study of omeprazole did not show ncreased tumor occurrence, but the study was not conclusive.

Esomeprazole was negative in the Ames mutation test, in the in vivo rat bone marrow cell chromosome aberration test, and the in vivo mouse nicronucleus test. Esomeprazole, however, was positive in the *in vitro* human lymphocyte chromosome aberration test. Omeprazole was lositive in the *in vitro* human lymphocyte chromosome aberration test, the *in vivo* mouse bone marrow cell chromosome aberration test, and

The potential effects of esomeprazole on fertility and reproductive performance were assessed using omeprazole studies. Omeprazole at oral doses up to 138 mg/kg/day in rats (about 34 times the human dose of 40 mg/day on a body surface area basis) was found to have no effect on

reproductive performance of parental animals. 13.2 Animal Toxicology and/or Pharmacology

Reproduction Studies
Reproduction studies have been performed in rats at oral doses up to 280 mg/kg/day (about 68 times an oral human dose of 40 mg on a body surface area basis) and in rabbits at oral doses up to 86 mg/kg/day (about 42 times an oral human dose of 40 mg on a body surface area basis) and have revealed no evidence of impaired fertility or harm to the fetus due to esomeprazole [see Use in Specific Populations (8.1)].

A 28-day toxicity study with a 14-day recovery phase was conducted in juvenile rats with esomeprazole magnesium at doses of 70 to 280 mg, kg/day (about 17 to 68 times a daily oral human dose of 40 mg on a body surface area basis). An increase in the number of deaths at the high dose of 280 mg/kg/day was observed when juvenile rats were administered esomeprazole magnesium from postnatal day 7 through postnatal day 35. In addition, doses equal to or greater than 140 mg/kg/day (about 34 times a daily oral human dose of 40 mg on a body surface area basis), produced treatment-related decreases in body weight (approximately 14%) and body eight gain, decreases in femur weight and femur length, and affected overall growth. Comparable findings described above have also been observed in this study with another esomeprazole salt, prazole strontium, at equimolar doses of esomeprazole.

14 CLINICAL STUDIES

14.1 Healing of EE in Adults

The healing rates of esomeprazole magnesium delayed-release capsules 40 mg, esomeprazole magnesium delayed-release capsules 20 mg, and omeprazole delayed-release capsules 20 mg (the approved dose for this indication) once daily were evaluated in adult patients with endoscopically diagnosed EE in four multicenter, double-blind, randomized studies. The healing rates at Weeks 4 and 8 were evaluated and are shown in Table 11:

Table 11: EE Healing Rate (Life-Table Analysis) in Adults with EE Treated with Esomeprazole Magnesium Delayed-Release Capsules or Omeprazole Delayed-Release Capsules Once Daily in Four Clinical Studies

	No. of Dotionto	Treatment Crown	EE Heali	ng Rates	Significance Level ⁷
udy	No. of Patients	Treatment Group	Week 4	Week 8	Significance Level
1	588	Esomeprazole magnesium delayed-release capsules 20 mg	68.7%	90.6%	N.S.
1	588	Omeprazole 20 mg	69.5%	88.3%	
	654	Esomeprazole magnesium delayed-release capsules 40 mg	75.9%	94.1%	p < 0.001
2	656	Esomeprazole magnesium delayed-release capsules 20 mg	70.5%	89.9%	p < 0.05
	650	Omeprazole 20 mg	64.7%	86.9%	
3	576	Esomeprazole magnesium delayed-release capsules 40 mg	71.5%	92.2%	N.S.
J	572	Omeprazole 20 mg	68.6%	89.8%	
4	1216	Esomeprazole magnesium delayed-release capsules 40 mg	81.7%	93.7%	p < 0.001
4	1209	Omeprazole 20 mg	68.7%	84.2%	
n-rar	k test vs. omeoraz	role 20 mg			

N.S.= not significant (p > 0.05)

In these same studies of patients with EE, sustained heartburn resolution and time to sustained heartburn resolution were evaluated and are

Table 12: Sustained Resolution1 of Heartburn in Adults with EE Treated with Esomeprazole Magnesium Delayed-Release Capsules or Omeprazole Delayed-Release Capsules Once Daily in Four Clinical Studies

			Cumulative Percent ² with Sustained Resolution		
Study	No. of Patients	Treatment Group	Day 14	Day 28	Significance Level ³
1	573	Esomeprazole magnesium 20 mg	64.3%	72.7%	N.S.
	555	Omeprazole 20 mg	64.1%	70.9%	
2	621	Esomeprazole magnesium 40 mg	64.8%	74.2%	p <0.001
	620	Esomeprazole magnesium 20 mg	62.9%	70.1%	N.S.
	626	Omeprazole 20 mg	56.5%	66.6%	
3	568	Esomeprazole magnesium 40 mg	65.4%	73.9%	N.S.
	551	Omeprazole 20 mg	65.5%	73.1%	
4	1187	Esomeprazole magnesium 40 mg	67.6%	75.1%	p <0.001
	1188	Omeprazole 20 mg	62.5%	70.8%	

2. Defined as the cumulative proportion of patients who have reached the start of sustained resolution 3. log-rank test vs. omeprazole 20 mg.

N.S. = not significant (p> 0.05)

In these four studies, the range of median days to the start of sustained resolution (defined as 7 consecutive days with no heartburn) was 5 days for esomeprazole magnesium 40 mg, 7 to 8 days for esomeprazole magnesium 20 mg and 7 to 9 days for omeprazole 20 mg. There are no comparisons of 40 mg of esomeprazole magnesium with 40 mg of omeprazole in clinical trials assessing either healing or symptomatic relief of EE.

14.2 Maintenance of Healing of EE in Adults

pulticenter, randomized, double-blind placebo-controlled 4-arm studies were conducted in adult patients with endoscopically confirmed. healed EE to evaluate esome prazole magnesium 40 mg (n=174), 20 mg (n=180), 10 mg (n=168) or placebo (n=171) once daily over six months

No additional clinical benefit was seen with esomeprazole magnesium 40 mg over esomeprazole magnesium 20 mg. Esomeprazole magnesium delayed-release capsules 40 mg once daily is not a recommended regimen for the maintenance of healing of EE in adults.

The percentages of patients that maintained healing of EE at the various time points are shown in the Figures 2 and 3: Figure 2: Maintenance of Healing Rates of EE in Adults by Month (Study 177)





Figure 3: Maintenance of EE Healing Rates in Adults by Month (Study 178)



Patients remained in remission significantly longer and the number of recurrences of EE was significantly less in patients treated with esomeprazole magnesium compared to placebo.

In both studies, the proportion of patients on esomeprazole magnesium who remained in remission and were free of heartburn and other GERD symptoms was well differentiated from placebo. In a third multicenter open label study of 808 patients treated for 12 months with esomeprazole magnesium 40 mg, the percentage of patients

that maintained healing of EE was 93.7% for six months and 89.4% for one year.

14.3 Symptomatic GERD in Adults

Two multicenter, randomized, double-blind, placebo-controlled studies were conducted in a total of 717 adult patients comparing four weeks of treatment with esomeprazole magnesium delayed-release capsules 20 mg or 40 mg once daily versus placebo for resolution of GERD symptoms. Patients had at least a 6-month history of heartburn episodes, no EE by endoscopy, and heartburn on at least four of the seven days

The percentage of patients that were symptom-free of heartburn was significantly higher in the esomeprazole magnesium groups compared to placebo at all follow-up visits (Weeks 1, 2, and 4).

No additional clinical benefit was seen with esomeprazole magnesium 40 mg over esomeprazole magnesium 20 mg. Esomeprazole magnesium 40 mg once daily is not a recommended regimen for the treatment of symptomatic GERD in adults. The percent of patients symptom-free of heartburn by day are shown in the Figures 4 and 5

Figure 4: Percent of Patients Symptom-Free of Heartburn by Day (Study 225)





Figure 5: Percent of Patients Symptom-Free of Heartburn by Day (Study 226)



In three European symptomatic GERD trials, esomeprazole magnesium 20 mg and 40 mg and omeprazole 20 mg were evaluated. No significant treatment related differences were seen.

14.4 Pediatric GERD

12 Years to 17 Years of Age
In a multicenter, randomized, double-blind, parallel-group study, 149 adolescent patients (12 to 17 years of age; 89 female; 124 Caucasian, 15
Black, 10 Other) with clinically diagnosed GERD were treated with esomeprazole magnesium delayed-release capsules 20 mg or 40 mg once daily for up to 8 weeks to evaluate safety and tolerability. Patients were not endoscopically characterized as to the presence or absence of EE.

14.5 Risk Reduction of NSAID-Associated Gastric Ulcer
Two multicenter, double-blind, placebo-controlled studies were conducted in adult patients at risk of developing gastric and/or duodenal ulcers associated with continuous use of non-selective and COX-2 selective NSAIDs. A total of 1429 patients were randomized across the 2 studies. Patients ranged in age from 19 to 89 (median age 66 years) with 71% female, 29% male; 83% Caucasian, 5% Black, 4% Asian, and 8% Others. Tations larged in Age from 19 de Original age Objects) with 17 or finale, 29 or finale, 30 or occurrent, 30 or loads, 30 or occurrent. At baseline, the patients in these studies were endoscopically confirmed not to have ulcers but were determined to be at risk for ulcer occurrence due to their age (at least 60 years) and/or history of a documented gastric or duodenal ulcer within the past 5 years. Patients receiving NSAID and treated with esomeprazole magnesium delayed-release capsules 20 mg or 40 mg once daily experienced significant reduction in gastric ulcer occurrences relative to placebo treatment at 26 weeks. See Table 13. No additional benefit was seen with esomeprazole magnesium 40 mg over esomeprazole magnesium 20 mg. Esomeprazole magnesium 40 mg once daily is not recommended regimen for the risk reduction of NSAID-associated gastric ulcer in adults. These studies did not demonstrate significant reduction in the development of NSAID-associated duodenal ulcer due to the low incidence.

Table 13: Cumulative Percentage of Patients at Least 60 Years of Age Taking NSAIDS Without Gastric Ulcers at 26 Weeks in Two Randomized

	Study	No. of Patients	Treatment Group	% of Patients Remaining Gastric Ulcer Free ¹		
	1	191 194 184	Esomeprazole magnesium delayed-release capsules 20 mg Esomeprazole magnesium delayed-release capsules 40 mg Placebo	95.4 96.7 88.2		
	2	267 271 257	Esomeprazole magnesium delayed-release capsules 20 mg Esomeprazole magnesium delayed-release capsules 40 mg Placebo	94.7 95.3 83.3		
1. %= Life Table Estimate. Significant difference from placebo (p<0.01).						

14.6 *H. pylori* Eradication in Adult Patients with Duodenal Ulcer Disease
Two multicenter, randomized, double-blind studies were conducted in adult patients using a 10-day treatment regimen of triple therapy (esomeprazole magnesium, amoxicillin and clarithromycin). The first study (191) compared esomeprazole magnesium delayed-release capsules 40 mg once daily in combination with amoxicillin 1000 mg twice daily and clarithromycin 500 mg twice daily to esomeprazole magnesium delayed-release capsules 40 mg once daily plus clarithromycin 500 mg twice daily. The second study (193) compared esomeprazole magnesium delayed-release capsules 40 mg once daily in combination with amoxicillin 1000 mg twice daily and clarithromycin 500 mg twice daily to esomeprazole magnesium 40 mg once daily. *H. pylori* eradication rates, defined as at least two negative tests and no positive tests from CLOtest®, histology and/or culture, at 4 weeks post-therapy were significantly higher in the esomeprazole magnesium, amoxicillin and clarithromycin group than in the esomeprazole magnesium and clarithromycin group or the esomeprazole magnesium delayed-release capsules alone group. The results are shown in Table 14:

Table 14: H. pylori Eradication Rates at 4 Weeks after 10 Day Treatment Regimen % of Adult Patients Cured [95% Confidence Interval]

Study	Treatment Group	Per-Protocol ¹	Intent-to-Treat ²
191	Esomeprazole magnesium, amoxicillin and clarithromycin	84% ³ [78, 89] (n=196)	77%³ [71, 82] (n=233)
	Esomeprazole magnesium and clarithromycin	55% [48, 62] (n=187)	52% [45, 59] (n=215)
193	Esomeprazole magnesium, amoxicillin and clarithromycin	85% ⁴ [74, 93] (n=67)	78% ⁴ [67, 87] (n=74)
	Esomeprazole magnesium	5% [0, 23] (n=22)	4% [0, 21] (n=24)

1. Patients were included in the analysis if they had *H. pylori* infection documented at baseline, had at least one endoscopically verified duodenal ulcer ≥ 0.5 cm in diameter at baseline or had a documented history of duodenal ulcer disease within the past 5 years, and were not

protocol violators. Patients who dropped out of the study due to an adverse reaction related to the study drug were included in the analysis Patients were included in the analysis if they had documented H. pylori infection at baseline, had at least one documented duodenal ulcer at baseline, or had a documented history of duodenal ulcer disease, and took at least one dose of study medication. All dropouts were included

as not H, pylori eradicated.

3 p < 0.05 compared to esomeprazole magnesium plus clarithromycin.

4 p < 0.05 compared to esomeprazole magnesium alone.

The percentage of patients with a healed baseline duodenal ulcer by 4 weeks after the 10-day treatment regimen in the esomeprazole magnesiur delayed-release capsules, amoxicillin and clarithromycin group was 75% (n=156) and 57% (n=60) respectively, in the 191 and 193 studies (per-protocol analysis).

14.7 Pathological Hypersecretory Conditions, Including Zollinger-Ellison Syndrome, in Adults
In a multicenter, open-label dose-escalation study of 21 adult patients (15 males and 6 females, 18 Caucasian and 3 Black, mean age of 56 years)
with pathological hypersecretory conditions, such as Zollinger-Ellison Syndrome, esomeprazole magnesium significantly inhibited gastric acid
secretion. The initial dosage of esomeprazole magnesium delayed-release capsules was 40 mg twice daily in 19 patients and 80 mg twice daily
in 2 patients. Total daily doses ranging from 80 mg to 240 mg for 12 months maintained gastric acid output below the target levels of 10 mEq/h in patients without prior gastric acid-reducing surgery and below 5 mEq/hr in patients with prior gastric acid-reducing surgery. At the Month 12 final visit, 18/20 (90%) patients had Basal Acid Output (BAO) under satisfactory control (median BAO = 0.17 mmol/hr). Of the 18 patients

with a starting dose of esomeprazole magnesium 40 mg twice daily, 13 (72%) had their BAO controlled with the original do

Table	Table 15: Adequate Acid Suppression at Final Visit by Dosage Regimen in Adult Patients with Pathological Hypersecretory Condition				
	Esomeprazole Magnesium dose at the Month 12 visit	BAO under adequate control at the Month 12 visit (N=20) ¹			
	40 mg twice daily	13/15			
	80 mg twice daily	4/4			
	80 ma three times daily	1/1			

16 HOW SUPPLIED/STORAGE AND HANDLING ium delayed-release capsules USP, 20 mg are White opaque size '4' hard gelatin capsule imprinted with "H" on cap and Esomeprazole magnesium delayed-release capsules U 'E2' on body filled with off white to pale yellow pellets.

regimen at the final visit. See Table 15.

NDC 31722-664-30 bottles of 30 NDC 31722-664-90 bottles of 90 NDC 31722-664-10 bottles of 1000

Esomeprazole magnesium delayed-release capsules USP, 40 mg are white opaque size '3' hard gelatin capsule imprinted with "H" on cap and 'E3' on body filled with off white to pale yellow pellets. NDC 31722-665-30 bottles of 30

NDC 31722-665-90 bottles of 90

1. One patient was not evaluated

NDC 31722-665-10 bottles of 1000 Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Keep esomeprazole magnesium delayed-release capsules, USP

Dispense in a tight container if the esomeprazole magnesium delayed-release capsules, USP product package is subdivided

Advise the patient to read the FDA-approved patient labeling (Medication Guide and Instructions for Use)

Acute Tubulointerstitial Nephritis
Advise the patient or caregiver to call the patient's healthcare provider immediately if they experience signs and/or symptoms associated with suspected acute TIN [see Warnings and Precautions (5.2)]. Clostridium difficile-Associated Diarrhea
Advise the patient or caregiver to immediately call the patient's healthcare provider if they experience diarrhea that does not improve [see Warnings]

Bone Fracture
Advise the patient or caregiver to report any fractures, especially of the hip, wrist or spine, to the patient's healthcare provider [see Warnings and Precautions (5.4)].

Advise the patient or caregiver to discontinue esomeprazole magnesium delayed-release capsules and immediately call the patient's healthcare provider for at first appearance of a severe cutaneous adverse reaction or other sign of hypersensitivity signs or symptoms associated with Severe Cutaneous Adverse Reactions [see Warnings and Precautions (5.5)].

<u>Cutaneous and Systemic Lupus Erythematosus</u>
Advise the patient or caregiver to immediately call the patient's healthcare provider for any new or worsening of symptoms associated with cutaneous or systemic lupus erythematosus [see Warnings and Precautions (5.6)]. Cyanocobalamin (Vitamin B-12) Deficiency.

Advise the patient or caregiver to report any clinical symptoms that may be associated with cyano

provider if they have been receiving esomeprazole magnesium delayed-release capsules for longer than 3 years [see Warnings and Precautions (5.8)].

Hypomagnesemia and Mineral Metabolism Advise the patient or caregiver to report any clinical symptoms that may be associated with hypomagnesemia, hypocalcemia, and/or hypokalemia to the patient's healthcare provider, if they have been receiving esomeprazole magnesium delayed-release capsules for at least 3 months [see Warnings and Precautions (5.9)].

Advise the patient or caregiver to report to their healthcare provider if starting treatment with rilpivirine-containing products, clopi John's Wort or rifampin; or, if they take high-dose methotrexate [see Contraindications (4), Warnings and Precautions (5.7, 5.10, 5.12)].

Antacids may be used concomitantly with esomeprazole magnesium delayed-release capsules Swallow esomeprazole magnesium delayed-release capsules whole; do not chew or crush the capsules. For patients who have difficulty swallowing capsules, esomeprazole magnesium delayed-release capsules can be opened, and the contents sprinkled on applesauce. Use with other foods is not recommended.

1. Add one tablespoon of applesauce to an empty bowl. The applesauce used should not be hot and should be soft enough to be

Administer the mixture immediately. Do not chew or crush the granules. Discard any remaining mixture. Do not store the mixture for future use

Open the esomeprazole magnesium delayed-release capsule and carefully empty the granules inside the capsule onto the

neprazole magnesium delayed-release capsules can also be administered via a nasogastric tube, as described in the Instructions for Use.

CAMBER Manufactured for: Camber Pharmaceuticals, Inc. Piscataway, NJ 08854

Revised: 04/2024

Hetero Labs Limited Jeedimetla, Hyderabad - 500 055,

swallowed without chewing.

the lulos acry ontal hells and

your doctor effects to F effects to F v should I sules? Store esc

ademarks of . The makers Hetero or its

the the tube

hake the syringe well for in the pand check for granul for they have dissolved or I

60 mt only a delaye

The (NG

owners are not