

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use DEFERASIROX ORAL GRANULES safely an effectively. See full prescribing information for DEFERASIROX ORAL GRANULES. DEFERASIROX oral granules, for oral use

Initial U.S. Approval: 2005

WARNING: RENAL FAILURE, HEPATIC FAILURE, and GASTROINTESTINAL HEMORRHAGE

Deferasirox may cause serious and fatal: acute kidney injury, including acute renal failure requiring dialysis and renal tubular toxicity including

hepatic toxicity, including failure (5.2)

Deferasirox therapy requires close patient monitoring, including laboratory tests of renal and hepatic

--INDICATIONS AND USAGE Deferasirox oral granules are an iron chelator indicated for the treatment of chronic iron overload due to blood transfusions

Deferasirox oral granules are indicated for the treatment of chronic iron overload in patients 10 years of ane and older with lent thalassemia (NTDT) syndromes, and with a liver iron (Fe) concentration (LIC) of at least 5 mg Fe

per gram of dry weight (Fe/g dw) and a serum ferritin greater than 300 mcg/L. (1.2) Limitations of Use: The safety and efficacy of deferasirox oral granules when administered with other iron chelation therapy have not been

---DOSAGE AND ADMINISTRATION----

Transfusional Iron Overload: Initial dose for patients with estimated glomerular filtration rate (eGFR) greater than $60\,\text{mL/min}/1.73\,\text{m}^2$ is $14\,\text{mg}$ per kg (calculated to nearest whole sachet content for granules) once daily. (2.1) NTDT Syndromes: Initial dose for patients with eGFR greater than 60 mL/min/1.73 m² is 7 mg per kg (calculated to

nearest whole sachet content for granules) once daily. (2.2) impairment, (2.1, 2.2, 2.3, 2.4)

---DOSAGE FORMS AND STRENGTHS---Granules: 90 mg, 180 mg, 360 mg. (3)

-CONTRAINDICATIONS-Estimated GFR less than 40 mL/min/1.73 m². (4)

Patients with poor performance status. (4)

Patients with high-risk myelodysplastic syndrome (MDS) (4)

Patients with advanced malignancies. (4)
Patients with platelet counts less than 50 x 10°/L. (4)

FULL PRESCRIBING INFORMATION: CONTENTS* WARNING: RENAL FAILURE, HEPATIC FAILURE, and GASTROINTESTINAL HEMORRHAGE

INDICATIONS AND USAGE $Treatment of Chronic Iron Overload \ Due to \ Blood \ Transfusions \ (Transfusional Iron \ Overload)$ Treatment of Chronic Iron Overload in Non-Transfusion-Dependent Thalassemia Syndromes

Acute Kidney Injury, Including Acute Renal Failure Requiring Dialysis and Renal Tubular Toxicity Including

DOSAGE AND ADMINISTRATION

Iron Overload in Non-Transfusion-Dependent Thalassemia Syndromes Use in Patients with Baseline Hepatic or Renal Impairment

Dose Modifications for Decreases in Renal Function While on Deferasirox oral Granules Dose Modifications Based on Concomitant Medications 3 DOSAGE FORMS AND STRENGTHS

CONTRAINDICATIONS WARNINGS AND PRECAUTIONS

Hepatic Toxicity and Failure

5.3 Gastrointestinal (GI) Ulceration, Hemorrhage, and Perforation Bone Marrow Sunn

Age-Related Risk of Toxicity 5.6 Overchelation

5.8 Severe Skin Reactions

5.9 Skin Rash 5.10 Auditory and Ocular Ab

ADVERSE REACTIONS Clinical Trials Experience

FULL PRESCRIBING INFORMATION

WARNING: RENAL FAILURE, HEPATIC FAILURE, and GASTROINTESTINAL HEMORRHAGE Deferasirox can cause acute renal failure and death, particularly in patients with comorbidities and

those who are in the advanced stages of their hematologic disorders. Evaluate baseline renal function prior to starting or increasing deferasirox dosing in all pati Deferasirox is contraindicated in adult and pediatric patients with eGFR less than 40 mL/min/1.73 m Measure serum creatinine in duplicate prior to initiation of therapy. Monitor renal function at least monthly. For patients with baseline renal impairment or increased risk of acute renal failure, monitor

renal function weekly for the first month, then at least monthly. Reduce the starting dose in patients with prexisting renal disease. During therapy, increase the frequency of monitoring and modify the dose for patients with an increased risk of renal impairment, including use of concomitant nephrotoxic drugs, and pediatric patients with volume depletion or overchelation (see Dosage and Administration (2.1, 2.4, 2.5), Warnings and Precautions (5.1), Adverse Reactions (6.1, 6.2)].

Measure serum transaminases and bilirubin in all patients prior to initiating treatment, every 2 weeks during the first month, and at least monthly th

Avoid use of deferasirox in patients with severe (Child-Pugh C) hepatic impairment and reduce the dose in patients with moderate (Child-Pugh B) hepatic impairment [see Dosage and Administration (2.4],

itestinal (GI) hemorrhages, which may be fatal, especially in elderly Deferasirox can cause gast patients who have advanced hematologic malignancies and/or low platelet cou

r patients and discontinue deferasirox for suspected GI ulceration or hemorrhage /see Warnings INDICATIONS AND USAGE

Deferasirox oral granules are indicated for the treatment of chronic iron overload due to blood transfusions (trans hemosiderosis) in patients 2 years of age and older. 1.2 Treatment of Chronic Iron Overload in Non-Transfusion-Dependent Thalassemia Syndrom Deferasing or all granules are indicated for the treatment of chronic iron overload in patients 10 years of age and older with non-transfusion-dependent thalassemia (NTDT) syndromes and with a liver iron concentration (LIC) of at least 5 milliograms

1.1 Treatment of Chronic Iron Overload Due to Blood Transfusions (Transfusional Iron Overload)

The safety and efficacy of deferasirox oral granules when administered with other iron chelation therapy have not been

2 DOSAGE AND ADMINISTRATION

2.1 Transfusional Iron Overload Deferacions therapy should only be considered when a natient has evidence of chronic transfusional iron overload. The evidence should include the transfusion of at least 100 mt /kg of packed red blood cells (e.g., at least 20 units of packed red blood cells for a 40 kg person or more in individuals weighing more than 40 kg), and a serum ferritin consistently greater than 1,000 mcg/L.

Prior to starting therapy, or increasing dose, evaluate: Baseline renal function

Obtain serum creatinine in duplicate (due to variations in measurements).

Calculate the estimated glomerular filtration rate (eGFR). Use a prediction equation appropriate for adult patients (e.g., CKD-EPI, MDRD method) and in pediatric patients (e.g., Schwartz equations). Obtain urinalyses and serum electrolytes to evaluate renal tubular function (see Dosage and Administration

(2.4), Warnings and Precautions (5.1)]. Serum transaminases and bilirubin [see Dosage and Administration (2.4), Warnings and Precautions (5.2)] Baseline auditory and ophthalmic examinations (see Warnings and Precautions (5.10))

Initiating Therapy: The recommended initial dose of deferasirox oral granules for patients 2 years of age and older with eGFR greater than 60 mL/min/1.73 m² is 14 mg per kg body weight orally, once daily. Calculate doses (mg per kg per day) to the nearest whole sachet content for granules. Changes in weight of pediatric patients over time must be taken into account when calculating

During Therapy: Monitor serum ferritin monthly and adjust the dose of deferasirox, if necessary, every 3 to 6 months based on serum

Use the minimum effective dose to achieve a trend of decreasing ferriting Make dose adjustments in steps of 3.5 or 7 mg per kg and tailor adjustments to the individual patient's response and

In patients not adequately controlled with doses of 21 mg per kg (e.g., serum ferritin levels persistently above 2,500 mcg/L and not showing a decreasing trend over time), doses of up to 28 mg per kg may be considered. Doses above 28 mg per kg are not recommended/see Warnings and Precautions (5.6!).

Adjust dose based on serum ferritin levels If the serum ferritin falls below 1,000 mcg/L at 2 consecutive visits, consider dose reduction especially if the deferasirox dose is greater than 17.5 mg/kg/day /see Adverse Reactions (6.1)].

If the serum ferritin falls below 500 mcg/L, interrupt deferasirox therapy to mini and continue monthly monitoring (see Warnings and Precautions (5.6)).

Evaluate the need for ongoing chelation therapy for patients whose conditions no longer require regular blood Use the minimum effective dose to maintain iron burden in the target range [see Warnings and Precautions

Monitor blood counts, liver function, renal function and ferritin monthly [see Warnings and Precautions (5.1, 5.2, 5.4]]. Interrupt deferasirox oral granules for pediatric patients who have acute illnesses, which can cause volume depletion, such as vomiting, diarrhea, or prolonged decreased oral intake, and monitor more frequently. Resume therapy as appropriate, based on assessments of renal function, when oral intake and volume status are normal

Dosage and Administration (2.4, 2.5), Warnings and Precautions (5.1), Use in Specific Populations (8.4), Clinical Pharmacology (12.3)].

 ${\bf 2.2} \quad Iron\,Overload\,in\,Non-Transfusion-Dependent\,Thalassemia\,Syndromes$ Deferasirox therapy should only be considered when a patient with NTDT syndrome has an LIC of at least 5 mg Fe/g dw and a serum ferritin greater than 300 mcg/L.

Prior to starting therapy, obtain: LIC by liver biopsy or by an FDA-cleared or approved method for identifying patients for treatment with deferasirox Serum ferritin level on at least 2 measurements 1-month apart [see Clinical Studies (14)]

Baseline renal function: Obtain serum creatinine in duplicate (due to variations in n Calculate the estimated glomerular filtration rate (eGFR). Use a prediction equation appropriate for adult

Baseline auditory and ophthalmic examinations (see Warnings and Precautions (5.10))

patients (e.g., CKD-EPI, MDRD method) and in pediatric patients (e.g., Schwartz equation Obtain urinalyses and serum electrolytes to evaluate renal tubular function [see Dosage and Administration (2.4), Warnings and Precautions (5.1)]. Serum transaminases and bilirubin [see Dosage and Administration (2.4), Warnings and Precautions (5.2)]

Initiating Therapy: nded initial dose of deferasirox for patients with eGFR greater than 60 mL/min/1.73 m² is 7 mg per kg body weight orally once daily. Calculate doses (mg per kg per day) to the nearest whole sachet content for granules

If the baseline LIC is greater than 15 mg Fe/g dw, consider increasing the dose to 14 mg/kg/day after 4 weeks.

During Therapy: itor serum ferritin monthly to assess the patient's response to therapy and to minimize the risk of overchelation (see Warnings and Precautions (5.6)). Interrupt treatment when serum ferritin is less than 300 mcg/L and obtain an

LIC to determine whether the LIC has fallen to less than 3 mg Fe/g dw. Use the minimum effective dose to achieve a trend of decreasing fe

After 6 months of therapy, if the LIC remains greater than 7 mg Fe/g dw, increase the dose of deferasirox to a

maximum of 14 mg/kg/day. Do not exceed a maximum of 14 mg/kg/day.

If after 6 months of therapy, the LIC is 3 to 7 mg Fe/g dw, continue treatment with deferasirox at no more than

Monitor blood counts, liver function, renal function and ferritin monthly (see Warnings and Precautions (5.1, 5.2,

 $Increase\ monitoring\ frequency\ for\ pediatric\ patients\ who\ have\ acute\ illness,\ which\ can\ cause\ volume\ depletion,\ such$ as vomiting, diarrhea, or prolonged decreased oral intake. Consider dose interruption until oral intake and volume status are normal [see Dosage and Administration (2.4, 2.5), Warnings and Precautions (5.1), Use in Specific

Populations (8.4), Clinical Pharmacology (12.3)]. Restart treatment when the LIC rises again to more than 5 mg Fe/g dw

Take deferasirox oral granules on an empty stomach or with a light meal (contains less than 7% fat content and nately 250 calories). Examples of light meals include 1 whole wheat English muffin, 1 packet jelly (0.5 ounces), and skim milk (8 fluid ounces) or a turkey sandwich (2 oz. turkey on whole wheat bread w/ lettuce, tomato, and 1 packet mustard/ [see Clinical Pharmacology (12.3]). Administer deferasino oral granules by sprinkling the full dose on soft food (e.g., yogurt or applesauce) immediately prior to use and administered orally. Deferasino oral granules should be taken once a day, preferably at the same time each day. Do not take deferasirox oral granules with aluminum-containing antacid For patients who are currently on chelation therapy with deferasirox tablets for oral suspension and converting to

deferasirox oral granules, the dose should be about 30% lower, rounded to the nearest whole sachet content for granules. $The table below \stackrel{-}{provides} additional information on dosing conversion to defer a sirox or all granules.$

	for oral suspension	Deferasirox Oral Granules
Transfusion-Dependent Iron Overload		
Starting Dose	20 mg/kg/day	14 mg/kg/day
Titration Increments	5 to 10 mg/kg	3.5 to 7 mg/kg
Maximum Dose	40 mg/kg/day	28 mg/kg/day
Non-Transfusion-Dependent Thalasser	nia Syndromes	
Starting Dose	10 mg/kg/day	7 mg/kg/day

----WARNINGS AND PRECAUTIONS--

Acute Kidney Injury: Measure serum creatinine in duplicate before starting therapy. Monitor renal function during

Fatal and Nonfatal Gastrointestinal (GI) Bleeding, Ulceration, and Irritation: Risk may be greater in patients who are aking deferasirox in combination with drugs that have known ulcerogenic or hemorrhagic potential. (5.3)

Hepatic Toxicity: Monitor hepatic function. Reduce dose or interrupt therapy for toxicity. (5.2)

Bone Marrow Suppression: Neutropenia, agranulocytosis, worsening anemia, and thrombocytopenia, including fatal events; monitor blood counts during deferasirox therapy. Interrupt therapy for toxicity. (5.4) Age-related Risk of Toxicity: Monitor elderly and pediatric patients closely for toxicity. (5.5)

Hypersensitivity Reactions: Discontinue deferasirox for severe reactions and institute medical intervention. (5.7) Severe Skin Reactions including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN), and drug reaction with eosinophilia and systemic symptoms (DRESS): Discontinue deferasirox. (5.8) -----ADVERSE REACTIONS----

In patients with transfusional iron overload, the most frequently occurring (greater than 5%) adverse reactions are diarrhea, vomiting, nausea, abdominal pain, skin rashes, and increases in serum creatinine. In deferasirox-treated patients with NTDT syndromes, the most frequently occurring (greater than 5%) adverse reactions are diarrhea, rash, and nausea. (6.1) To report SUSPECTED ADVERSE REACTIONS, contact Annora Pharma Private Limited at 1-866-495-1995 or

FDA at 1-800-FDA-1088 or www.fda.gov/medwatch. -----DRUG INTERACTIONS---

Do not take deferasirox with aluminum-containing antacid preparations. (7.1)
Deferasirox increases the exposure of repaglinide. Consider repaglinide dose reduction and monitor blood glucose Avoid the use of deferasirox with theophylline as theophylline levels could be increased. (7.4)

Deferasirox increases exposure of busulfan. Monitor plasma concentrations of busulfan when coadministered with deferasirox to allow dose adjustment of busulfan, as needed. (7.7)

....USE IN SPECIFIC POPULATIONS Lactation: Advise women not to breastfeed. (8.2) See 17 for PATIENT COUNSELING INFORMATION and Medication Guide

DRUG INTERACTIONS Aluminum-Containing Antacid Preparations

7.2 Agents Metabolized by CYP3A4

Agents Metabolized by CYP1A2 7.5 Agents Inducing UDP-glucuronosyltransferase (UGT) Metabolism Bile Acid Sequestrants

7.7 Busulfan 8 USE IN SPECIFIC POPULATIONS

Pregnancy 8.2 Lactation

8.3 Females and Males of Reproductive Potential 8.5 Geriatric Use

8.7 Hepatic Impairment

10 OVERDOSAGE DESCRIPTION

CLINICAL PHARMACOLOGY 12.1 Mechanism of Action

12.2 Pharmacodynamics 12.3 Pharmacokinetics 13 NONCLINICAL TOXICOLOGY

14 CLINICAL STUDIES 16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

		Deferasirox Tablets for oral suspension	Deferasirox Oral Granules	
	Titration Increments	5 to 10 mg/kg	3.5 to 7 mg/kg	
	Maximum Dose	20 mg/kg/day	14 mg/kg/day	

nts with Baseline Hepatic Impairm Mild (Child-Pugh A) Hepatic Impairment: No dose adjustment is necessary

Moderate (Child-Pugh B) Hepatic Impairment: Reduce the starting dose by 50%. Severe (Child-Pugh C) Hepatic Impairment: Avoid deferasirox oral granules [see Warnings and Precautions (5.2), Use in Specific Populations (8.7)].

Patients with Baseline Renal Impair Do not use deferasirox in adult or pediatric patients with eGFR less than 40 mL/min/1.73 m² (see Dosage and Administration

For patients with renal impairment (eGFR 40 to 60 mL/min/1.73 m²), reduce the starting dose by 50% [see Use in Specific Exercise caution in pediatric patients with eGFR between 40 and 60 mL/minute/1.73 m^2 . If treatment is needed, use the

2.5 Dose Modifications for Decreases in Renal Function While on Deferasirox Oral Granules $Deferasirox\ or al\ granules\ are\ contraindicated\ in\ patients\ with\ eGFR\ less\ than\ 40\ mL/min/1.73\ m^2/see\ Contraindications\ (4)/s.$ For decreases in renal function while receiving deferasirox oral granules (see Warnings and Precautions (5.1)), modify the

Transfusional Iron Overload

If the serum creatinine increases by 33% or more above the average baseline measurement, repeat the serum creatinine within 1 week, and if still elevated by 33% or more, reduce the dose by 7 mg per kg. Pediatric Patients (ages 2 years to 17 years): Reduce the dose by 7 mg per kg if eGFR decreases by greater than 33% below the average baseline me

Interrupt deferasirox oral granules for acute illnesses, which can cause volume depletion, such as vomiting, diarrhea,

or prolonged decreased oral intake, and monitor more frequently. Resume therapy as appropriate, based on assessments of renal function, when oral intake and volume status are normal. Avoid use of other nephrotoxic drugs (see Warnings and Precautions (5.1)). In the setting of decreased renal function, evaluate the risk benefit profile of continued deferasirox oral granules use Use the minimum effective deferasirox oral granules dose and monitor renal function more frequently, by evaluating tubular and glomerular function. Titrate dosing based on renal injury. Consider dose reduction or interruption and less nephrotoxic-therapies until improvement of renal function. If signs of renal tubular or glomerular injury occur in the

presence of other risk factors such as volume depletion, reduce or interrupt deferasirox oral granules to prevent evere and irreversible renal injury (see Warnings and Precautions (5.1)). All Patients (regardless of age):

Discontinue therapy for eGFR less than 40 mL/min/1.73 m²/see Contraindications (4)].

creatining within 1 week, and if still elevated by 33% or more, interrupt therapy if the dose is 3.5 mg per kg, or reduce by 50% if the dose is 7 or 14 mg per kg. $Reduce\ the\ dose\ by\ 3.5\ mg\ per\ kg\ if\ eGFR\ decreases\ by\ greater\ than\ 33\%\ below\ the\ average\ baseline\ measurement$

If the serum creatinine increases by 33% or more above the average baseline measurement, repeat the serum

Pediatric Patients (ages 10 years to 17 years

Increase monitoring frequency for pediatric patients who have acute illnesses, which can cause volume depletion such as vomiting, diarrhea, or prolonged decreased oral intake. Consider dose interruption until oral intake and

volume status are normal. Avoid use of other nephrotoxic drugs (see Warnings and Precautions (5.1)). In the setting of decreased renal function, evaluate the risk benefit profile of continued deferasirox oral granules use.

Use the minimum effective deferasirox oral granules dose and monitor renal function more frequently, by evaluating tubular and glomerular function. Titrate dosing based on renal injury. Consider dose reduction or interruption and less therapies until improvement of renal function. If signs of renal tubular or glomerular injury occur in the presence of other risk factors such as volume depletion, reduce or interrupt deferasirox oral granules to prevent severe and irreversible renal injury (see Warnings and Precautions (5.1)).

All Patients (regardless of age): Discontinue therapy for eGFR less than 40 mL/min/1.73 m²/see Contraindications (4)/.

2.6 Dose Modifications Based on Concomitant Medications UDP-glucuronosyltransferases (UGT) Inducers

Concomitant use of UGT inducers decreases systemic exposure. Avoid the concomitant use of strong UGT inducers (e.g., rifampicin, phenytoin, phenobarbital, ritonavir). If you must administer deferasirox oral granules with a strong UGT inducer onsider increasing the initial dose by 50%, and monitor serum ferritin levels and clinical responses for further dose nodification/see Dosage and Administration (2.1, 2.2), Drug Interactions (7.5).

Concomitant use of bile acid sequestrants decreases systemic exposure. Avoid the concomitant use of bile acid sequestrants (e.g., cholestyramine, colesevelam, colestipol). If you must administer deferasirox oral granules with a bille acid sequestrant, consider increasing the initial dose by 50%, and monitor serum ferritin levels and clinical responses for further dose modification/see Dosage and Administration (2.1, 2.2), Drug Interactions (7.6)].

DOSAGE FORMS AND STRENGTHS

90 mg Deferasirox oral granules supplied in cartons containing 30 child resistant foil sachets. Each sachet contains 162 mg of white to off white granular powder, equivalent to 90 mg deferasirox. 180 mg Deferasirox oral granules supplied in cartons containing 30 child resistant foil sachets. Each sachet contain

 $360\,mg$ Deferasirox oral granules supplied in cartons containing 30 child resistant foil sachets. Each sachet contains $648\,mg$ of white to off white granular powder, equivalent to $360\,mg$ deferasirox. asinot una grainuse a Contrainucioscum patients with.

Estimated GFR liess than 40 m.[min] 1.73 m²[see Dosage and Administration (2.5), Warnings and Precautions (5.1)];

Poor performance status (see Warnings and Precautions (5.1, 5.3));

324 mg of white to off white granular powder, equivalent to 180 mg deferasirox.

High-risk myelodysplastic syndromes (this patient population was not studied and is not expected to benefit from Advanced malignancies (see Warnings and Precautions (5.1, 5.3));

Platelet counts less than 50 x 10°/L [see Warnings and Precautions (5.3, 5.4]]; Known hypersensitivity to deferasirox or any component of deferasirox oral granules (see Warnings and Precautions

Acute Kidney Injury, Including Acute Renal Failure Requiring Dialysis and Renal Tubular Toxicity Including sirox is contraindicated in patients with eGFR less than 40 mL/min/1.73 m². Exercise caution in pediatric patients with eGFR between 40 and 60 mL/min/1.73 m². If treatment is needed, use the minimum effective dose and monitor rena function frequently. Individualize dose titration based on improvement in renal injury [see Use in Specific Populations (8.6]]. For patients with renal impairment (eGFR 40 to 60 mL/min/1.73 m²) reduce the starting dose by 50% (see Dosage and

Administration (2.4, 2.5), Use in Specific Populations (8.6)]. Deferasirox can cause acute kidney injury including renal failure requiring dialysis that has resulted in fatal outcomes. Based on postmarketing experience, most fatalities have occurred in patients with multiple comorbidities and who were in nced stages of their hematological disorders. In the clinical trials, adults and pediatric deferasirox-treated patients with no preexisting renal disease experienced dose-dependent mild, non-progressive increases in serum creat inuria. Preexisting renal disease and concomitant use of other nephrotoxic drugs may increase the risk of acute kidne injury in adult and pediatric patients. Acute illnesses associated with volume depletion and overchelation may increase the risk of acute kidney injury in pediatric patients. In pediatric patients, small decreases in eGFR can result in increases in deferasirox exposure, particularly in younger patients with body surface area typical of patients less than age 7 years. This can lead to a cycle of worsening renal function and further increases in deferasirox tablets for oral suspension exposure, unless the dose is reduced or interrupted. Renal tubular toxicity, including acquired Fanconi syndrome, has been reported in patients treated with deferasirox, most commonly in pediatric patients with beta-thalassemia and serum ferritin levels le than 1,500 mcg/L (see Warnings and Precautions (5.6), Adverse Reactions (6.1, 6.2), Use in Specific Populations (8.4) Clinical Pharmacology (12.3)].

validated for use in adult and pediatric patients to estimate GFR. Obtain serum electrolytes and urinalysis in all patients to evaluate renal tubular function (see Dosage and Administration (2.1, 2.2)). Monitor all patients for changes in eGFR and for renal tubular toxicity weekly during the first month after initiation o modification of therapy and at least monthly thereafter. Dose reduction or interruption may be considered if abnormalities occur in levels of markers of renal tubular function and/or as clinically indicated. Monitor serum ferritin monthly to evaluate

Evaluate renal glomerular and tubular function before initiating therapy or increasing the dose. Use prediction equations

for overchelation. Use the minimum dose to establish and maintain a low iron burden. Monitor renal function more frequently in patients with preexisting renal disease or decreased renal function. In pediatric patients, interrupt deferasirox during acute illnesses, which can cause volume depletion such as vomiting, diarrhea, or prolonged decreased oral intake, and monitor renal function more frequently. Promptly correct fluid deficits to prevent renal injury. Resume therapy as appropriate based on assessments of renal function, when oral intake and volume status are normal (see Dosage and ration (2.5), Warnings and Precautions (5.6), Adverse Reactions (6.1, 6.2), Use in Specific Popu 5.2 Hepatic Toxicity and Failure irox can cause hepatic injury, fatal in some patients. In Study 1, 4 patients (1,3%) discontinued deferasirox because

toxicity appears to be more common in patients greater than 55 years of age. Hepatic failure was more common in patients with significant comorbidities, including liver cirrhosis and multiorgan failure [see Adverse Reactions (6.1)]. Acute liver injury and failure, including fatal outcomes, have occurred in pediatric defensions-treated natients. Liver failure occurred in association with acute kidney injury in pediatric patients at risk for overchelation during a volume-deple event. Interrupt deferasirox therapy when acute liver injury or acute kidney injury is suspected and during volume depletion

iron burden (see Dosage and Administration (2.5), Warnings and Precautions (5.6), Adverse Reactions (6.1)]. Measure transaminases (aspartate transaminase (AST) and alanine transaminase (ALT)) and bilirubin in all patients befor the initiation of treatment and every 2 weeks during the first month and at least monthly thereafter. Consider dose ons or interruption of treatment for severe or persistent elevations. Avoid the use of deferasirox in patients with severe (Child-Pugh C) hepatic impairment. Reduce the starting dose in patients

with moderate (Child-Pugh B) hepatic impairment (see Dosage and Administration (2.4), Use in Specific Populations (8.7)].

Monitor liver and renal function more frequently in pediatric patients who are receiving deferasirox in the 14 to 28 mg/kg/day range and when iron burden is approaching normal. Use the minimum effective dose to achieve and maintain a low

Patients with mild (Child-Pugh A) or moderate (Child-Pugh B) hepatic impairment may be at higher risk for hepatic toxicity. ${\bf 5.3} \quad {\bf Gastrointestinal \, (GI) \, Ulceration, Hemorrhage, and \, Perforation}$

GI hemorrhage, including deaths, has been reported, especially in elderly patients who had advanced hematologi malignancies and/or low platelet counts. Nonfatal upper Gl irritation, ulceration and hemorrhage have been reported in patients, including children and adolescents, receiving deferasirox /see Adverse Reactions (6.1)/. Monitor for signs and symptoms of GI ulceration and hemorrhage during deferasirox therapy, and promptly initiate additional evaluation and symptoms of or discretion and reinformage during deterasions, therapy, and promptly illustre adultional evaluation at treatment if a serious GI adverse reaction is suspected. The risk of GI hemorrhage may be increased when administering deferasiros in combination with drugs that have ulcerogenic or hemorrhagic potential, such as nonsteroidal anti-inflammatory drugs (NSAIDs), corticosteroids, oral bisphosphonates, or anticoagulants. There have been reports of ulcers complicated with GI perforation (including fatal outcome) /see Adverse Reactions (6.2)/

5.4 Bone Marrow Suppression nia, agranulocytosis, worsening anemia, and thrombocytopenia, including fatal events, have been reported i patients treated with deferasirox. Preexisting hematologic disorders may increase this risk. Monitor blood counts in all patients. Interrupt treatment with deferasirox in patients who develop cytopenias until the cause of the cytopenia has been determined. Deferasirox is contraindicated in patients with platelet counts below 50 x 10°/L.

predominantly in elderly patients. Monitor elderly patients treated with deferasirox more frequently for toxicity [see Use in Specific Populations (8.5)

Pediatric Patients Deferasirox has been associated with serious and fatal adverse reactions in pediatric patients in the postmarketing setting. These events were frequently associated with volume depletion or with continued deferasirox tablets for oral suspension doses in the 20 to 40 mg/kg/day range equivalent to 14 to 28 mg/kg/day deferasirox when body iron burden was approaching or in the normal range. Interrupt deferasirox in patients with volume depletion, and resume deferasirox when renal function and fluid volume have normalized. Monitor liver and renal function more frequently during volume depletion and in patients receiving deferance in a commence of the comme Precautions (5.6), Use in Specific Populations (8.4)].

5.6 Overchelation For patients with transfusional iron overload, measure serum ferritin monthly to assess the patient's response to therapy and minimize the risk of overchelation. An analysis of pediatric patients treated with deferasirox tablets for oral suspension in p clinical trials (n = 158), found a higher rate of renal adverse reactions among patients receiving doses greater than $25\,\mathrm{mg/kg/day}$ equivalent to 17.5 mg/kg/day deferasirox while their serum ferritin values were less than 1,000 mcg/L. Consider dose reduction or closer monitoring of renal and hepatic function, and serum ferritin levels during these periods. Use the minimum effective dose to maintain a low-iron burden [see Adverse Reactions (6.1), Use in Specific Populations (8.4)].

dose is greater than 17.5 mg/kg/day (see Adverse Reactions (6.1)). If the serum ferritin falls below 500 mcg/L, interrup therapy with deferasirox and continue monthly monitoring. Evaluate the need for ongoing chelation for patients whose conditions do not require regular blood transfusions. Use the minimum effective dose to maintain iron burden in the target range. Continued administration of deferasirox in the 14 to 28 mg/kg/day range, when the body iron burden is approaching of hin the normal range can result in life-threatening adverse reactions (see Dosage and Administration (2.1)]. For patients with NTDT, measure LIC by liver biopsy or by using an FDA-cleared or approved method for monitoring patient receiving deferasirox therapy every 6 months on treatment. Interrupt deferasirox adm

If the serum ferritin falls below 1,000 mcg/L at 2 consecutive visits, consider dose reduction, especially if the deferasiron

onfirmatory LIC /see Clinical Studies (14) Deferacions may cause serious hypersensitivity reactions (such as anaphylaxis and angioedema), with the onset of the reaction usually occurring within the first month of treatment (see Adverse Reactions (6.21). If reactions are severe, discontinue deferasirox and institute appropriate medical intervention. Deferasirox is contraindicated in patients with known hypersensitivity to deferasirox products and should not be reintroduced in patients who have experienced previous

mg Fe/g dw. Measure serum ferritin monthly, and if the serum ferritin falls below 300 mcg/L, interrupt deferasirox and

hypersensitivity reactions on deferasirox products due to the risk of anaphylactic shock ous adverse reactions (SCARs), including Stevens-Johnson syndrome (SJS), toxic epidermal necrolysis (TEN) and drug reaction with eosinophilia and systemic symptoms (DRESS) which could be life-threatening or fatal have been reported during deferasirox therapy [see Adverse Reactions (6.1, 6.2)]. Cases of erythema multiforme have been observed.

Advise patients of the signs and symptoms of severe skin reactions, and closely monitor. If any severe skin reactions are suspected, discontinue deferasirox immediately and do not reintroduce deferasirox therapy. Rashes may occur during deferasirox treatment [see Adverse Reactions (6.1)]. For rashes of mild to moderate severity,

deferasirox may be continued without dose adjustment, since the rash often resolves spontaneously. In severe cases, interrupt treatment with deferasirox. Reintroduction at a lower dose with escalation may be considered after resolution o Auditory disturbances (high frequency hearing loss, decreased hearing), and ocular disturbances (lens opacities, cataracts, elevations in intraocular pressure, and retinal disorders) were reported at a frequency of less than 1% with deferasirox

received deferasirox tablets for oral suspension doses greater than 25 mg/kg/day equivalent to 17.5 mg/kg/day deferasiro. when serum ferritin was less than 1,000 mcg/L/see Warnings and Precautions (5.6), Perform auditory and ophthalmic testing (including slit lamp examinations and dilated fundoscopy) before starting deferasirox treatment and thereafter at regular intervals (every 12 months). If disturbances are noted, monitor mor

therapy in the clinical studies. The frequency of auditory adverse reactions was increased among pediatric patients, who

frequently. Consider dose reduction or inter ADVERSE REACTIONS

The following clinically significant adverse reactions are also discussed in other sections of the labeling: Acute Kidney Injury, Including Acute Renal Failure Requiring Dialysis, and Renal Tubular Toxicity Including Fanconi

Syndrome [see Warnings and Precautions (5.1, 5.6)] Hepatic Toxicity and Failure (see Warnings and Precautions (5.2, 5.6)) GI Hemorrhage (see Warnings and Precautions (5.3)) Bone Marrow Suppression [see Warnings and Precautions (5.4]]

ere Skin Reactions (see Warnings and Precautions (5.8)) Skin Rash (see Warnings and Precautions (5.9)) Auditory and Ocular Abnormalities [see Warnings and Precautions (5.10)]

Hypersensitivity (see Warnings and Precautions (5.7))

practice. Deferasirox was evaluated in healthy volunteer trials. Currently, there are no clinical data in patients with deferasirox oral granules. Deferasirox oral granules contain the same active ingredient as deferasirox tablets for oral suspension. The following adverse reactions have been reported with deferasirox tablets for oral suspensio A total of 700 adult and pediatric patients were treated with deferasirox for 48 weeks in premarketing studies. These

45% were male, 70% were Caucasian, and 292 patients were less than 16 years of age. In the sickle cell disease population,

a, 99 with rare anemias, and 132 with sickle cell dis

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of

83% of patients were black. Median treatment duration among the sickle cell patients was 51 weeks. Of the 700 patients treated, 469 (403 beta-thalassemia and 66 rare anemias) were entered into extensions of the original clinical protocols. In ongoing extension studies, median durations of treatment were 88 to 205 weeks. Six hundred twenty-seven (627) patients with myelodysplastic syndrome (MDS) were enrolled across 5 uncontrolled trials. These studies varied in duration from 1 to 5 years. The discontinuation rate across studies in the first year was 46% Adverse Events (AEs) 20%, withdrawal of consent 10%, death 8%, other 4%, lab abnormalities 3%, and lack of efficacy 1%). Among 47 patients enrolled in the study of 5-year duration, 10 remained on deferasirox at the completion of the study. Table 1 displays adverse reactions occurring in greater than 5% of deferasirox-treated beta-thalassemia patients (Study 1), sickle cell disease patients (Study 3), and patients with MDS (MDS pool). Abdominal pain, nausea, vomiting, diarrhea, skin

rashes, and increases in serum creatinine were the most frequent adverse reactions reported with a suspected to deferasirox. Gastrointestinal symptoms, increases in serum creatinine, and skin rash were dose related Table 1. Adverse Reactions Occurring in > 5% of Deferasirox-treated Patients in Study 1, Study 3, and MDS

		udy 1 alassemia)	Study 3 (Sickle Cell Disease)		MDS Pool	
Adverse Reactions	Deferasirox N = 296 n (%)	Deferoxamine N = 290 n (%)	Deferasirox N = 132 n (%)	Deferoxamine N = 63 n (%)	Deferasirox N = 627 n (%)	
Abdominal Pain ^b	63 (21)	41 (14)	37 (28)	9 (14)	145 (23)	
Diarrhea	35 (12)	21 (7)	26 (20)	3 (5)	297 (47)	
Creatinine Increased ^c	33 (11)	0 (0)	9 (7)	0	89 (14)	
Nausea	31 (11)	14 (5)	30 (23)	7 (11)	161 (26)	
Vomiting	30 (10)	28 (10)	28 (21)	10 (16)	83 (13)	
Rash	25 (8)	9 (3)	14 (11)	3 (5)	83 (13)	

In Study 1, a total of 113 (38%) patients treated with deferasirox had increases in serum creatining greater than 33% above seline on 2 separate occasions (Table 2) and 25 (8%) patients required dose reductions. Increases in se appeared to be dose related (see Warnings and Precautions (5.1)]. In this study, 17 (6%) patients treated with deferasiron developed elevations in serum glutamic-pyruvic transaminase (SGPT)/ALT levels greater than 5 times the upper limit of normal (ULN) at 2 consecutive visits. Of these, 2 patients had liver biopsy proven drug-induced hepatitis and both discontinued deferasirox therapy (see Warnings and Precautions (5.2)). An additional 2 patients, who did not have elevations in SGPT/ALT greater than 5 times the ULN, discontinued defeasions because of increased SGPT/ALT. Increases in transaminases did not appear to be dose related. Adverse reactions that led to discontinuations included abnormal liver

function tests (2 patients) and drug-induced hepatitis (2 patients), skin rash, glycosuria/proteinuria, Henoch Schönlein

In Study 3, a total of 48 (36%) patients treated with deferasirox had increases in serum creatining greater than 33% above

baseline on 2 separate occasions (Table 2) [see Warnings and Precautions (5.1]]. Of the patients who experienced creatining

increases in Study 3, 8 deferasirox-treated patients required dose reductions. In this study, 5 patients in the deferasirox

*Adverse reaction frequencies are based on AEs reported regardless of relationship to study drug.

mnia, drug fever, and cataract (1 patient each).

group developed elevations in SGPT/ALT levels greater than 5 times the ULN at 2 consecutive visits and 1 patient subsequently had deferasirox permanently discontinued. Four additional patients discontinued due to adverse reactions with a suspected relationship to study drug, including diarrhea, pancreatitis associated with gallstones, atypica In the MDS pool, in the first year, a total of 229 (37%) patients treated with deferasirox had increases in serum creatining greater than 33% above baseline on 2 consecutive occasions (Table 2) and 8 (3.5%) patients permanently disco Warnings and Precautions (5.1). A total of 5 (0.8%) patients developed SGPT/ALT levels greater than 5 times the ULN at 2 diarrhea, nausea, rash, and vomiting. Death was reported in the first year in 52 (8%) of patients [see Clinical Studies (14)].

Table 2. Number (%) of Patients with Increases in Serum Creatinine or SGPT/ALT in Study 1, Study 3, and MDS

	Study 1 (Beta-thalassemia)		Study 3 (Sickle Cell Disease)		MDS Pool	
Laboratory Parameter	Deferasirox N = 296 n (%)	Deferoxamine N = 290 n (%)	Deferasirox N = 132 n (%)	Deferoxamine N = 63 n (%)	Deferasirox N = 627 n (%)	
Serum Creatinine						
Creatinine increase > 33% at 2 consecutive post-baseline visits	113 (38)	41 (14)	48 (36)	14 (22)	229 (37)	
Creatinine increase > 33% and > ULN at 2 consecutive post-baseline visits	7 (2)	1 (0)	3 (2)	2 (3)	126 (20)	
SGPT/ALT						
SGPT/ALT > 5 x ULN at 2 post- baseline visits	25 (8)	7 (2)	2 (2)	0	9 (1)	
SGPT/ALT > 5 x ULN at 2 consecutive post-baseline visits	17 (6)	5 (2)	5 (4)	0	5 (1)	

Non-Transfusion-Dependent Thalassemia Syndromes In Study 5, 110 patients with NTDT received 1 year of treatment with deferasirox 5 or 10 mg/kg/day and 56 patients received placebo in a double-blind, randomized trial. In Study 6, 130 of the patients who completed Study 5 were treated with open-label deferasirox at 5, 10, or 20 mg/kg/day (depending on the baseline LIC) for 1 year (see Clinical Studies (14)). In Study 7, 134 patients with NTDT of 10 years of age or older with iron overload, received deferasirox tablets for oral suspension for up to 5 years, at a starting dose of 10 mg/kg/day followed by dose adjustment at Week 4, and then approximately every 6 months thereafter based on LIC levels. Table 3 and 4 display the frequency of adverse reactions in

Abbreviations: ALT, alanine transaminase; MDS, myelodysplastic syndrome; SGPT, serum glutamic-pyruvic

at \geq 5% of patients in Study 5.

transaminase; ULN, upper limit of normal.

	Study 5		Study 6	Study 7
	Deferasirox N = 110 n (%)	Placebo N = 56 n (%)	Deferasirox N = 130 n (%)	Deferasirox N = 134 n (%)
Any adverse reaction	31 (28)	9 (16)	27 (21)	50 (37)
Nausea	7 (6)	4 (7)	2 (2)	7 (5)
Rash	7 (6)	1 (2)	2 (2)	3 (2) ^a
Diarrhea	5 (5)	1 (2)	7 (5)	8 (6)

additional adverse reactions with a suspected relationship to study drug occurring in > 5% of patients in Study 6 In Study 5, 1 patient in the placebo 10 mg/kg/day group experienced an ALT increase to greater than 5 times ULN and greater than 2 times baseline (Table 4). Three deferasirox-treated patients (all in the 10 mg/kg/day group) had 2 consecutiv n creatinine level increases greater than 33% from baseline and greater than ULN. Serum creat in all 3 patients (in 1 spontaneously and in the other 2 after drug interruption). Two additional cases of ALT increase and

*The occurrence of nausea, and rash are included for Study 6 and rash for Study 7 for consistency. There were n

2 additional cases of serum creatinine increase were observed in the 1-year extension of Study 5. The number (%) o atients with NTDT with increase in serum creatinine or SGPT/ALT in Study 5, Study 6 and Study 7 are presented in Table 4

	Stu	dy 5	Study 6	Study 7	
Laboratory Parameter	Deferasirox N = 110 n (%)	Placebo N = 56 n (%)	Deferasirox N = 130 n (%)	Deferasirox N = 134 n (%)	
Serum creatinine (> 33% increase from baseline and > ULN at ≥ 2 consecutive post-baseline values)	3 (3)	0	2 (2)	2 (2)	
SGPT/ALT (> 5 x ULN and > 2 x baseline)	1 (1)	1 (2)	2 (2)	1 (1)	

In clinical studies, urine protein was measured monthly, Intermittent proteinuria (urine protein/creatinine ratio greater that

0.6 mg/mg) occurred in 18.6% of deferasirox-treated patients compared to 7.2% of deferoxamine-treated patients in Study
1/see Warnings and Precautions (5.1).

Deferasirox (dee fer' a sir ox) **Oral Granules**

MEDICATION GUIDE

What is the most important information I should know about deferasirox oral granules?

Deferasirox oral granules can cause serious side effects, including: Kidney problems: Deferasirox oral granules can cause sudden (acute) kidney problems, including kidney failure that may require treatment with dialysis, and may cause death. Deaths have happened mostly in people who also have other health problems and had a blood disorder that was in an advanced stage. Adults and children who already have kidney problems and are taking certain medicines with deferasirox oral granules may also have an increased risk of sudden kidney problems. Be sure to tell your healthcare provider about all the medicines you take during treatment with deferasirox oral granules.

Your healthcare provider should do blood and urine tests to check your or your child's kidney function before and during treatment with deferasirox oral granules. Call your healthcare provider right away if:

 your child becomes sick with fever, vomiting, or diarrhea and cannot drink fluids normally during treatment with deferasirox oral granules. Your child may be dehydrated. Your healthcare provider may need to temporarily stop treatment with deferasirox oral granules and treat your child for dehydration to help prevent kidney problems. Your healthcare provider may monitor your child's kidney function more closely.

 you notice that you or your child are passing less urine than usual during treatment with deferasirox oral granules.

Liver problems. Deferasirox oral granules can cause liver problems, including liver failure that can sometimes cause death. Liver problems with deferasirox oral granules may be more common in people who are over 55 years of age but can also happen in children. Liver failure has happened more often in people with cirrhosis of the liver and failure of other organs. Liver failure has also happened along with kidney problems in certain children who become dehydrated. See "Kidney problems" above.

Your healthcare provider should do blood tests to check your liver function before you start and regularly during treatment with deferasirox oral granules. Call your healthcare provider right away, if you develop any of the following signs and symptoms:

 drowsiness yellowing or increased

yellowing of your skin or eyes upper right stomach-area

Bleeding, ulcers, and tears of the stomach or intestine. Severe stomach and intestine bleeding (hemorrhage) that have caused death have happened in some people treated with deferasirox oral granules, especially in elderly people who have advanced blood cancers or low platelet counts. Some people have also had ulcers of the stomach or intestine, sometimes with tears (perforation) that have caused death. In some people who have taken deferasirox oral granules, including children and adolescents, irritation of the upper gastrointestinal tract, ulcers, and bleeding have happened, but did not cause death.

Your risk of severe bleeding (hemorrhage) may be increased if you take deferasirox oral granules along with other medicines that can cause ulcers or bleeding, such as:

drugs (NSAIDs) medicines called oral bisphosphonates blood thinner medicines corticosteroids

nonsteroidal anti-inflammatory • certain osteoporosis

your healthcare provider may stop deferasirox oral granules. Elderly people may be at a higher risk of developing serious side effects and death due to serious side effects with deferasirox oral granules. Your healthcare provider may need to monitor you more closely during

Before you start taking deferasirox oral granules, tell your healthcare provider if

you are taking one of these medicines. Ask your healthcare provider if you are

not sure. If you develop an ulcer of the stomach or intestine, or severe bleeding,

 Tell your healthcare provider if you get heartburn during treatment with deferasirox oral granules. Get emergency medical help right away if you vomit blood or pass

black or bloody stools, or if you have severe stomach-area

treatment with deferasirox oral granules.

What are deferasirox oral granules?

(abdomen) pain during treatment with deferasirox oral granules. See "What are the possible side effects of deferasirox oral granules?" for more information about side effects.

people 2 years of age and older who have an increased amount of iron in their blood for a long period of time (chronic), caused by repeated certain people 10 years of age or older with thalassemia who have an

Deferasirox oral granules is a prescription medicine that is used to treat:

increased amount of iron in their blood but who are not receiving regular blood transfusions It is not known if deferasirox oral granules are safe and effective when used

with other medicines to treat an increased amount of iron in the blood. It is not known if deferasirox oral granules are safe and effective for treating children under 2 years of age who have an increased amount of iron in their blood for a long period of time (chronic) caused by repeated

It is not known if deferasirox oral granules are safe and effective for treating children under 10 years of age with thalassemia who have an increased amount of iron in their blood, but who are not receiving regular blood transfusions.

Do not take deferasirox oral granules if you:

have high-risk myelodysplastic syndrome (MDS)

have certain kidney problems

have advanced cancer

have a low platelet count

are allergic to deferasirox or any of the ingredients in deferasirox oral granules. See the end of this medication guide for a list of the ingredients in deferasirox oral granules.

medical conditions listed above. Before taking deferasirox oral granules tell your healthcare

Ask your healthcare provider if you are not sure if you have any of the

provider about all of your medical conditions, including if you: have kidney problems

blood transfusions.

have liver problems have advanced cancer. See "Do not take deferasirox oral granules

have a blood disorder that may increase your risk for bleeding are pregnant or plan to become pregnant. It is not known if deferasirox oral granules can harm your unborn baby. Hormonal forms of birth control may not be as effective if used during treatment with deferasirox oral granules. You could become pregnant. Talk to your healthcare provider about other birth control options that you can use during this time. Tell your healthcare provider right away if you become pregnant during treatment with deferasirox oral granules.

are breastfeeding or plan to breastfeed. It is not known if deferasirox

passes into your breast milk and can harm your baby. You and your

healthcare provider should decide if you will take deferasirox oral granules or breastfeed. You should not do both. Tell your healthcare provider about all the medicines you take, including prescription medicines, over-the-counter medicines, vitamins, and herbal supplements. Some medicines may affect how deferasirox oral granules work, and deferasirox oral granules may affect how other

medicines work. Also, vour risk of sudden kidney problems or severe

bleeding may be increased if you take deferasirox oral granules

with certain medicines. See "What is the most important

 Avoid taking the following medicines during treatment with deferasirox oral granules:

information I should know about deferasirox oral granules?"

o antacid products (medicines used to treat heartburn) that contain

Artwork information				
Customer	Camber	Market	USA	
Dimensions (mm)	370 x 630 mm	Non Printing Colors	Die cut	
Pharma Code No.	Front-416 & Back-417			
Printing Colours (01)	Black			

Prepared by: Approved by: PD QA PK PK **RA**

- o certain medicines to lower your cholesterol, called bile acid

Ask your healthcare provider if you are not sure if you take one of these medicines.

Know the medicines you take. Keep a list of them to show your healthcare provider or pharmacist when you get a new medicine.

How should I take deferasirox oral granules?

- Take deferasirox oral granules exactly as your healthcare provider
- Do not change your dose of deferasirox oral granules or stop taking unless your healthcare provider tells you to.
- Deferasirox comes as oral granules.
- O You may take deferasirox oral granules on an empty stomach or with a light meal. Examples of a light meal include:
 - 1 whole wheat English muffin, 1 packet of jelly (0.5 ounce),
 - and skim milk (8 fluid ounces), or A turkey sandwich (2 ounces of turkey on whole wheat bread

with lettuce, tomato, and 1 packet of mustard)

- Taking deferasirox oral granules: o Sprinkle the prescribed dose of granules onto soft food such as
- yogurt or applesauce right before taking your dose.
- o Take the sprinkles and soft food by mouth. Do not take deferasirox oral granules with aluminum-containing
- antacid products. See "Before taking deferasirox oral granules". Tell your healthcare provider if you or your child gain or lose any weight. Your or your child's dose of deferasirox oral granules may
- need to be adjusted. If you take the diabetes medicine repaglinide during treatment with deferasirox oral granules, you may need to test your blood sugar (glucose) levels more often. Follow your healthcare provider's
- instructions about how often to test your blood sugar during this time. Your healthcare provider should do blood and urine tests before, and during treatment to check how you respond to deferasirox oral granules, and to monitor you for side effects. Your healthcare provider may change your dose, or temporarily or permanently stop deferasirox
- oral granules if you have certain side effects. In people who have thalassemia, your healthcare provider will check the amount of iron in your liver before and during treatment with deferasirox oral granules.
- If you or your child take too much deferasirox oral granules, call your healthcare provider right away or go to the nearest hospital emergency room. Symptoms that can happen if you take too much deferasirox oral granules include: stomach-area (abdominal) pain, diarrhea, nausea and vomiting.

What should I avoid while taking deferasirox oral granules?

Deferasirox oral granules may cause dizziness. Avoid driving or operating machinery until you know how deferasirox oral granules affect you. Do not drive or operate machinery if deferasirox oral granules make you dizzy.

What are the possible side effects of deferasirox oral granules? Deferasirox oral granules can cause serious side effects, including:

See "What is the most important information I should know

about deferasirox oral granules?"

- Effects on your bone marrow. Deferasirox oral granules can affect your bone marrow and cause you to have low white blood cell count which can be serious, decreased platelets, or worsening of your anemia, and may lead to death. Your risk for effects on your bone marrow may be increased if you already have other blood disorders. Your healthcare provider will do blood tests to monitor your blood cell counts for these problems.
- Serious allergic reactions. Deferasirox oral granules may cause serious allergic reactions, which usually start within the first month of treatment. **Get medical help right away if you develop any of the** following symptoms of a serious allergic reaction including:
- o difficulty in breathing or swallowing
- o swelling of the face, lips, mouth, tongue, or throat
- o chest pain o severe itching of the skin with a red rash or raised bumps
- o rapid heartbeat
- o hives
- o feeling faint
- Skin rash and severe skin reactions. Skin rashes are common with deferasirox oral granules. If you get a more severe rash, your healthcare provider may temporarily stop deferasirox oral granules. Severe skin reactions can also happen with deferasirox oral granules and can be life-threatening or lead to death. Get medical help right away if you develop any one or more of the following signs and symptoms of a severe skin reaction, including:
- o rash or red skin o skin peeling o blisters on your lips, o high fever or flu-like symptoms
- or around your mouth or eyes o enlarged lymph nodes o mouth sores
- Hearing and vision problems. Deferasirox oral granules can cause
- decreased hearing and changes in your vision including cataracts, increased pressure in your eye, and problems with your retinas. Your healthcare provider should do hearing and vision tests before you start and then regularly during treatment. Your healthcare provider may decrease your dose or stop deferasirox oral granules if you develop hearing or vision problems.

The most common side effects in anyone who takes deferasirox oral granules include: diarrhea and nausea.

Other common side effects in people with too much iron in their blood due to repeated blood transfusions include: vomiting, stomach-area (abdomen) pain, and an abnormal kidney function blood test.

These are not all the possible side effects of deferasirox oral granules. Call your doctor for medical advice about side effects. You may report side

effects to FDA at 1-800-FDA-1088. How should I store deferasirox oral granules?

- Store deferasirox oral granules at room temperature between 68°F to 77°F (20°C to 25°C).
- Deferasirox oral granules come in child-resistant sachets.
- Keep the bottle closed tightly and away from moisture.
- Keep deferasirox oral granules and all medicines out of the reach of children.

General information about the safe and effective use of deferasirox oral granules

Medicines are sometimes prescribed for purposes other than those listed in a Medication Guide. Do not use deferasirox oral granules for a condition for which it was not prescribed. Do not give it to other people, even if they have the same symptoms you have. It may harm them. You can ask your healthcare provider or pharmacist for more information about deferasirox oral granules.

What are the ingredients in deferasirox oral granules? Deferasirox oral granules:

Active ingredient: deferasirox

Inactive ingredients: colloidal silicon dioxide, crospovidone, microcrystalline cellulose, poloxamer (188), povidone (K30), and sodium stearyl fumarate.

For more information, call 1-866-495-1995 This Medication Guide has been approved by the U.S. Food and Drug

Administration.

Medication guide available at http://camberpharma.com/medication-guides

CAMBER Manufactured for: Camber Pharmaceuticals, Inc. Piscataway, NJ 08854 By: Annora Pharma Pvt. Ltd.

 $\ensuremath{\text{Sangareddy}}$ - 502313, Telangana, India.

In the population of more than 5,000 patients with transfusional iron overload, who have been treated with deferasirox during clinical trials, adverse reactions occurring in 0.1% to 1% of patients included gastritis, edema, sleep disorder, pigmentation disorder, dizziness, anxiety, maculopathy, cholelithiasis, pyrexia, fatigue, laryngeal pain, cataract, hearing loss, GI hemorrhage, gastric ulcer (including multiple ulcers), duodenal ulcer, renal tubular disorder (Fanconi syndrome), and acute pancreatitis (with and without underlying biliary conditions). Adverse reactions occurring in 0.01% to 0.1% of patients included optic neuritis, esophagitis, erythema multiforme, and drug reaction with eosinophilia and systemic symptoms (DRESS). Adverse reactions, which most frequently led to dose interruption or dose adjustment during clinical trials were rash, GI disorders, infections, increased serum creat

Pooled Analysis of Pediatric Clinical Trial Data

A nested case control analysis was conducted within a deferasirox tablets for oral suspension pediatric-pooled clinical trial dataset to evaluate the effects of dose and serum ferritin level, separately and combined, on kidney function. Among 1213 children (aged 2 to 15 years) with transfusion-dependent thalassemia, 162 cases of acute kidney injury (eGFR ≤ 90 mL/min/1.73 m²) and 621 matched-controls with normal kidney function (eGFR ≥ 120 mL/min/1.73 m²) were identified

- A 26% increased risk of acute kidney injury was observed with each 5 mg/kg increase in daily deferasirox tablets for oral suspension dosage equivalent to 3.5 mg/kg deferasirox, starting at 20 mg/kg/day equivalent to 14 mg/kg/day deferasirox (95% confidence interval (Cl): 1.08 to 1.48).
- A~25%~increased~risk~for~acute~kidney~injury~was~observed~with~each~250~mcg/L~decrease~in~serum~ferritin~starting~at
- Among pediatric patients with a serum ferritin < 1,000 mcg/L, those who received deferasirox tablets for oral suspension dosage > 30 mg/kg/day, equivalent to 21 mg/kg/day deferasirox compared to those who received lower dosages, had a higher risk for acute kidney injury (Odds ratio (OR) = 4.47, 95% Cl: 1.25 to 15.95), consistent with
- In addition, a cohort-based analysis of ARs was conducted in the deferasirox tablets for oral suspension pediatric pooled clinical trial data. Pediatric patients who received deferasirox tablets for oral suspension dose > 25 mg/kg/day equivalent to 17.5 mg/kg/day deferasirox when their serum ferritin was < 1,000 mcg/L (n = 158), had a 6-fold greater rate of renal adverse reactions (Incidence Rate Ratio (IRR) = 6.00, 95% CI: 1.75 to 21.36), and a 2-fold greater rate of dose interruptions IRRR = 2.06, 95% Cl: 1.33 to 3.17) compared to the time-period prior to meeting these simultaneous criteria. Adverse reactions of special interest (cytopenia, renal, hearing, and GI disorders) occurred 1.9-fold more frequently when these

Precautions (5.6)1. 6.2 Postmarketing Experience

The following adverse reactions have been spontaneously reported during post-approval use of deferasirox in the ransfusional iron overload setting. Because these reactions are reported voluntarily from a population of uncertain size, in which patients may have received concomitant medication, it is not always possible to reliably estimate frequency or stablish a causal relationship to drug exposure

simultaneous criteria were met, compared to preceding time-periods (IRR = 1.91, 95% CI: 1.05 to 3.48) /see Warnings and

Skin and Subcutaneous Tissue Disorders: Stevens-Johnson syndrome (SJS), leukocytoclastic vasculitis, urticaria, alopecia, Immune System Disorders: hypersensitivity reactions (including anaphylactic reaction and angioedema)

Hepatobiliary Disorders: hepatic failure GI Disorders: GI perforation

Blood and Lymphatic System Disorders: worsening anemia

5-Year Pediatric Registry 5-teaf Peudutic negastry
In a 5-year observational study, 267 pediatric patients 2 to < 6 years of age (at enrollment) with transfusional hemosiderosis received deferasirox. Of the 242 patients who had pre- and post-baseline eGFR measurements, 116 (48%)

Renal and Urinary Disorders: acute renal failure tuhulointerstitial nenhritis

natients had a decrease in eGFR of > 33% observed at least once. Twenty-one (18%) of these 116 natients with decreased eGFR had a dose interruption, and 15 (13%) of these 116 patients had a dose decrease within 30 days. Adverse reaction leading to permanent discontinuation from the study included liver injury (n = 11), vomiting (n = 2), renal tubular disorder (n = 1), proteinuria (n = 1), hematuria (n = 1), upper GI hemorrhage (n = 1), abdominal pain (n = 1), and hypokalemia (n = 1) 7 DRUG INTERACTIONS 7.1 Aluminum-Containing Antacid Preparations

The concomitant administration of deferasirox and aluminum-containing antacid preparations has not been formally studied

Although deferasirox has a lower affinity for aluminum than for iron, do not take deferasirox with aluminum-containing

Deferasirox may induce CYP3A4 resulting in a decrease in CYP3A4 substrate concentration when these drugs are coadministered. Closely monitor patients for signs of reduced effectiveness when deferasirox is administered with drugs metabolized by CYP3A4 (e.g., alfentanil, aprepitant, budesonide, buspirone, conivaptan, cyclosporine, darifenacin, darunavir, dasatinib, dihydroergotamine, dronedarone, eletriptan, eplerenone, ergotamine, everolimus, felodipine, fentanyl, hormonal contraceptive agents, indinavir, fluticasone, lopinavir, lovastatin, lurasidone, maraviroc, midazolam, nisoldipine, pimozide, quetiapine, quinidine, saquinavir, sildenafil, simvastatin, sirolimus, tacrolimus, tolvaptan, tipranavir, triazolam, ticanrelor. and vardenafil) (see Clinical Pharmacology (12.3)).

7.3 Agents Metabolized by CYP2C8

Deferasirox inhibits CYP2C8 resulting in an increase in CYP2C8 substrate (e.g., repaglinide and paclitaxel) concentration when these drugs are coadministered. If deferasirox and repaglinide are used concomitantly, consider decreasing the dose of repaglinide and perform careful monitoring of blood glucose levels. Closely monitor patients for signs of exposure related toxicity when deferasirox is coadministered with other CYP2C8 substrates [see Clinical Pharmacology (12.3]].

7.4 Agents Metabolized by CYP1A2

Deferasirox inhibits CYP1A2 resulting in an increase in CYP1A2 substrate (e.g., alosetron, caffeine, duloxetine, melatonin, ramelteon, tacrine, theophylline, tizanidine) concentration when these drugs are coadministered. An increase in theophylline ntrations could lead to clinically significant theophylline induced CNS or other adverse reactions. Avoid the concomitant use of theophylline or other CYP1A2 substrates with a narrow therapeutic index (e.g., tizanidine) with deferasirox. Monitor theophylline concentrations and consider theophylline dose modification if you must coadminister theophylline with deferasirox. Closely monitor patients for signs of exposure related toxicity when deferasirox is coadministered with other drugs metabolized by CYP1A2/see Clinical Pharmacology (12.3)].

7.5 Agents Inducing UDP-glucuronosyltransferase (UGT) Metabolism Deferasirox is a substrate of UGT1A1 and to a lesser extent UGT1A3. The concomitant use of deferasirox with strong UGT inducers (e.g., rifampicin, phenytoin, phenobarbital, ritonavir) may result in a decrease in deferasirox efficacy due to a possible decrease in deferasirox concentration. Avoid the concomitant use of strong UGT inducers with deferasirox. Consider increasing the initial dose of deferasirox if you must coadminister these agents together */see Dosage and* Administration (2.5), Clinical Pharmacology (12.3)].

7.6 Bile Acid Sequestrants Avoid the concomitant use of bile acid sequestrants (e.g., cholestyramine, colesevelam, colestipol) with deferasirox due to a possible decrease in deferasirox concentration. If you must coadminister these agents together, consider increasing the tial dose of deferasirox (see Dosage and Administration (2.5), Clinical Pharmacology (12.3)].

Increased exposure of busulfan was observed with concomitant use with deferasirox. Monitor plasma concentrations of busulfan when coadministered with deferasirox to allow dose adjustment of busulfan as needed *[see Clinical Pharmacology*

USE IN SPECIFIC POPULATIONS

There are no studies with the use of deferasirox in pregnant women to inform drug-associated risks. Administration of deferasirox to rats during pregnancy resulted in decreased offspring viability and an increase in renal anomalies in male offspring at doses that were about or less than the recommended human dose on a mg/m² basis. No fetal effects were noted in pregnant rabbits at doses equivalent to the human recommended dose on an mg/m² basis. Deferasirox

should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus. The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies had a background risk of hirth defect, loss, or other adverse outcomes. However, the background risk in the U.S. general population of major birth defects is 2% to 4% and of miscarriage is 15% to 20% of clinically recognized pregna

organogenesis at doses up to 100 mg/kg/day in rats and 50 mg/kg/day in rabbits (1.2 times the maximum rechuman dose (MRHD) on an mg/m²basis). These doses resulted in maternal toxicity but no fetal harm was observed. elopmental study, pregnant rats received oral deferasirox daily from organogenesis through lactation day 20 at doses of 10, 30, and 90 mg/kg/day (0.1, 0.3, and 1.0 times the MRHD on a mg/m² basis). Maternal toxicity, loss of litters, and decreased offspring viability occurred at 90 mg/kg/day (1.0 times the MRHD on a mg/m² basis), and increases in renal anomalies in male offspring occurred at 30 mg/kg/day (0.3 times the MRHD on a mg/m² basis). 8.2 Lactation

In embryo-fetal developmental studies, prepared rats and rabbits received oral deferasirox during the period of

No data are available regarding the presence of deferasirox or its metabolites in human milk, the effects of the drug on the Because many drugs are excreted in human milk and because of the potential for serious adverse reactions in a breastfeeding child from deferasirox and its metabolites, a decision should be made whether to discontinue breastfeeding or to discontinue the drug, taking into account the importance of the drug to the mother

8.3 Females and Males of Reproductive Potential

Counsel patients to use non-hormonal method(s) of contraception since deferasirox can render hormonal contraceptives 8.4 Pediatric Use

The safety and effectiveness of deferasirox have been established in pediatric patients 2 years of age and older for the treatment of transfusional iron overload (see Dosage and Administration (2.1)). Safety and effectiveness have not been established in pediatric patients less than 2 years of age for the treatment of

Pediatric approval for treatment of transfusional iron overload was based on clinical studies of 292 pediatric patients 2 years to less than 16 years of age with various congenital and acquired anemias. Seventy percent of these patients had beta-thalassemia [see Indications and Usage (11, Dosage and Administration (2.1), Clinical Studies (14)]. In those clinical studies, 173 children (ages 2 to < 12 years) and 119 adolescents (ages 12 to < 17 years) were exposed to deferasirox. A trial conducted in treatment naïve pediatric patients, 2 to < 18 years of age with transfusional iron overload (NCT02435212) did not provide additional relevant information about the safety or effectiveness of the deferasirox granules dosage form compared to the deferasirox oral tablets for suspension dosage form. ron Overload in Non-Transfusion-Dependent Thalassemia Syndromes

The safety and effectiveness of deferasirox have been established in patients 10 years of age and older for the treatment of chronic iron overload with non-transfusion-dependent thalassemia (NTDT) syndromes (see Dosage and Administration Safety and effectiveness have not been established in patients less than 10 years of age with chronic iron overload in NTDT

Pediatric approval for treatment of NTDT syndromes with liver iron (Fe) concentration (LIC) of at least 5 mg Fe per gram of dry weight and a serum ferritin greater than 300 mcg/L was based on 16 pediatric patients treated with deferasirox therapy (10 years to less than 16 years of age) with chronic iron overload and NTDT. Use of deferasirox in these age groups is Indications and Usage (1.2), Dosage and Administration (2.2), Clinical Studies (14)].

In general, risk factors for deferasirox-associated kidney injury include preexisting renal disease, volume depletion overchelation, and concomitant use of other nephrotoxic drugs. Acute kidney injury, and acute liver injury and failure has occurred in pediatric patients. In a pooled safety analysis, pediatric patients with higher deferasirox exposures had a greater probability of renal toxicity and decreased renal function, resulting in increased deferasirox exposure and progressive rena toxicity/kidney injury. Higher rates of renal AEs have been identified among pediatric patients receiving deferasirox tablets toxicity/numey injury. Injurie rates of relian Acs have user identified an injury general patients receiving user assists a user for oral suspension doses greater than 25 mg/kg/day edirects from the superior of the patients received by the patients of th Reactions (6.1, 6.2)].

 ${\it Monitoring recommendations for all pediatric patients with Transfusional Iron \it Overload and NTDT$ nded that serum ferritin be monitored every month to assess the patient's response to therapy and to minimiz the risk of overchelation [see Warnings and Precautions (5.6)].

Monitor renal function by estimating GFR using an eGFR prediction equation appropriate for pediatric patients and evaluate renal tubular function. Monitor renal function more frequently in pediatric patients in the presence of renal toxicity risk factors, including episodes of dehydration, fever and acute illness that may result in volume depletion or decreased renal perfusion. Use the minimum effective dose [see Warnings and Precautions (5.1)]. Interrupt deferasirox in pediatric patients with transfusional iron overload, and consider dose interruption in pediatric

patients with non-transfusion-dependent iron overload, for acute illnesses, which can cause volume depletion, such as vomiting, diarrhea, or prolonged decreased oral intake, and monitor more frequently. Resume therapy as appropriate, base on assessments of renal function, when oral intake and volume status are normal. Evaluate the risk benefit profile of continued deferasirox use in the setting of decreased renal function. Avoid use of other nephrotoxic drugs (see Dosage and Administration (2.5), Warnings and Precautions (5.1)). Juvenile Animal Toxicity Data Renal toxicity was observed in adult mice, rats, and marmoset monkeys administered deferasirox at therapeutic doses. In a

neonatal and juvenile toxicity study in rats, deferasirox was administered orally from postpartum Day 7 through 70, which equates to a human age range of term neonate through adolescence. Increased renal toxicity was identified in juvenile rats compared to adult rats at a dose based on mg/m2 approximately 0.4 times the recommended dose of 20 mg/kg/day. A higher

8.5 Geriatric Use Four hundred thirty-one (431) patients greater than or equal to 65 years of age were studied in clinical trials of deferasirox in the transfusional iron overload setting. Two hundred twenty-five (225) of these patients were between 65 and 75 years of age while 206 were greater than or equal to 75 years of age. The majority of these nationts had myelodysplastic syndrome (MDS) (n = 393). In these trials, elderly patients experienced a higher frequency of adverse reactions than younger patients Monitor elderly patients for early signs or symptoms of adverse reactions that may require a dose adjustment. Elderly nationts are at increased risk for toxicity due to the greater frequency of decreased benatic, renal, or cardiac function, and of tant disease or other drug therapy. Dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range.

In elderly patients, including those with MDS, individualize the decision to remove accumulated iron based on clinical circumstances and the anticipated clinical benefit and risks of deferasirox tablets for oral suspension therapy

with renal impairment (eGFR 40 to 60 mL/min/1.73 m2), reduce the starting dose by 50% (see Dosage and Administration $\textit{(2.4), Clinical Pharmacology (12.3)/}. Exercise caution in pediatric patients with an eGFR between 40 and 60 \, mL/min/1.73 \, m^{-2} \, m^{-2}$ (see Dosage and Administration (2.41). If treatment is needed, use the minimum effective dose with enhanced monitoring of glomerular and renal tubular function. Individualize dose titration based on improvement in renal injury [see Dosage and Administration (2.4, 2.5)].

Deferasirox can cause glomerular dysfunction, renal tubular toxicity, or both, and can result in acute renal failure. Monitor all patients closely for changes in eGFR and renal tubular dysfunction during deferasirox treatment. If either develops, consider dose reduction, interruption or discontinuation of deferasirox until glomerular or renal tubular function returns to baseline [see Dosage and Administration (2.4, 2.5), Warnings and Precautions (5.1)]. 8.7 Hepatic Impairment

Avoid use in patients with severe (Child-Pugh C) hepatic impairment. For patients with moderate (Child-Pugh B) hepatic impairment, reduce the starting dose by 50%. Closely monitor patients with mild (Child-Pugh A) or moderate (Child-Pugh B) hepatic impairment for efficacy and adverse reactions that may require dose titration (see Dosage and Administration (2.4). Warnings and Precautions (5.2), Clinical Pharmacology (12.3)]


Cases of overdose (2 to 3 times the prescribed dose for several weeks) have been reported. In one case, this resulted in

hepatitis which resolved without long-term consequences after a dose interruption. In one pediatric case, a dose of 2 to 3 times the prescribed dose for 6 days resulted in acute renal failure requiring hemofiltration and acute liver injury/failure, Stimes the prescribed uses for Outperformers and the state relating the provided for the prescribed with intensive care support. Single doses of deferasions up to 80 mg per kg per day with the tablet for oral suspension formulation in iron-overloaded beta-thalassemic patients have been tolerated with nausea and diarrhea noted. In healthy subjects, single doses of up to 40 mg per kg per day with the tablet for oral suspension formulation were Early signs of acute overdose are digestive effects such as abdominal pain, diarrhea, nausea, and vomiting. Hepatic and

ntinuation. An erroneously administered single dose of 90 mg/kg led to Fanconi syndrome which resolved after

There is no specific antidote for deferasirox. In case of overdose, it may be treated with induction of vomiting or gastrio lavage, and by symptomatic treatr 11 DESCRIPTION

Deferasirox is an iron-chelating agent provided as a oral granules for oral use. Deferasirox is designated chemically as [4-[(37.5E)-3.5-bis(6-oxo-1-cyclohexa-2.4-dienylidene)-1.2.4-triazolidin-1-yllbenzoic acid and has the following structural



Deferasirox is an off white to slightly yellow color crystalline powder. It has a molecular formula C2, H3, N3O4 and molecular weight of 373.36 g/mol. It is practically in-soluble in water and freely soluble in dimethyl formami Deferasirox oral granules contain 90 mg, 180 mg, or 360 mg deferasirox. Inactive ingredients include colloidal silicon dioxide,

(188), povidone (K30), sodium stearyl fumarate crystalline cellulose.

12 CLINICAL PHARMACOLOGY 12.1 Mechanism of Action

Deferasirox is an orally active chelator that is selective for iron (as Fe³*). It is a tridentate ligand that binds iron with high affinity in a 2:1 ratio. Although deferasirox has very low affinity for zinc and copper, there are variable decreases in the serum concentration of these trace metals after the administration of deferasirox. The clinical significance of thes

12.2 Pharmacodynamic that deferasirox (10, 20, and 40 mg per kg per day) was able to induce a mean net iron excretion (0.119, 0.329, and 0.445 within the clinically relevant range (0.1 to 0.5 mg per kg per day). Iro

An analysis of pooled pediatric clinical trial data found a statistically significant relationship between exposure and the probability of renal toxicity (increase in serum creatinine and urinary protein), resulting in a decrease in renal function Decreases in renal function resulted in an increase in deferasirox exposure which may increase the probability of renal

Cardiac Electrophysiolog At the maximum approved recommended dose, deferasirox does not prolong the QT interval to any clinically relevant extent 12.3 Pharmacokinetics

Based on studies in patients with the tablet for oral suspension, deferasirox is absorbed following oral administration with median times to maximum plasma concentration (T___) of about 1.5 to 4 hours. In healthy subjects, deferasions showed comparable T_{max}. The maximal concentrations (C_{max}) and area under the curve (AUC_{0 to 24} h, AUC_x) of deferasirox increase approximately linearly with dose after both single administration and under steady-state conditions. Exposure to deferasirox increased by an accumulation factor of 1.3 to 2.3 after multiple doses with the tablet for oral suspension formulation

The bioavailability (as measured by AUC,) of deferasirox oral granules was 52% greater than with deferasirox tablets for The unovariability last ineasure up you be least up to the defension or all guidelines was 32% greater timel with reflection or all suspension. After strength-adjustment, the mean AUC_in of the defensions or all granules (i.e., 4 x 90 mg strength) was similar to that of defension x tablets for oral suspension (i.e., 500 mg strength) under fasting conditions; however, the mean C..., was increased by 34%. The 34% increase in C..., observed with deferasirox oral granules is not clinically meaningful. The administration of deferasirox oral granules with a soft meal (e.g., yogurt and applesauce) or with a low-fat (approximately 450 calories with fat content approximately 30% of total calories) indicated that the AUC_{erf} and C_{max} after a low-fat meal or soft foods were similar to that under fasting conditions. The administration of deferasirox oral granules with

Deferasirox is highly (~99%) protein bound almost exclusively to serum albumin. The percentage of deferasirox confined to

high-fat meal (approximately 1,000 calories with fat content greater than 50% of total calories) increased $\overset{\circ}{AUC}_{int}$ by 18%

with no changes in C_{max} compared to that under fasting conditions (see Dosage and Administration (2.3)).

Glucuronidation is the main metabolic pathway for deferasirox, with subsequent biliary excretion. Deconjugation of glucuronidates in the intestine and subsequent reabsorption (enterohepatic recycling) is likely to occur. Deferasirox is mainly uronidated by UGT1A1 and to a lesser extent UGT1A3. CYP450-catalyzed (oxidative) metabolism of deferasiro appears to be minor in humans (about 8%). Deconjugation of glucuronide metabolites in the intestine and subsequent reabsorption (enterohepatic recycling) was confirmed in a healthy subjects study in which the administration of deferasirox resulted in a 45% decrease in deferasirox exposure (AUC,,) by interfering with the enterohepatic recycling of

Deferasirox and metabolites are primarily (84% of the dose) excreted in the feces. Renal excretion of deferasirox and metabolites is minimal (8% of the dose). The mean elimination half-life (t₁₂) ranged from 8 to 16 hours following oral

Midazolam: The concomitant administration of deferasirox tablets for oral suspension and CYP3A4 probe substrate midazolam resulted in a decrease of midazolam C_{max} by 23% and AUC_{at} by 17%. In the clinical setting, this effect may be more

pronounced, as the study was not adequately designed to conclusively assess the potential induction of CYP3A4 by Repaglinide: The concomitant administration of deferasirox tablets for oral suspension (30 mg per kg/day for 4 days) and the

CYP2C8 probe substrate repaglinide (single dose of 0.5 mg) increased repaglinide AUC, at 0.3-fold and C, as of 1.6-fold (see Drug Interactions (7.3)]. Theophylline: The concomitant administration of deferasirox tablets for oral suspension (repeated dose of 30 mg per kg/day)

and the CYP1A2 substrate theophylline (single dose of 120 mg) resulted in an approximate doubling of the theophylline AUC_{uu} and elimination half-life. The single dose C_{uu} was not affected, but an increase in theophylline C_{uu} is expected to occur with chronic dosing /see Drug Interactions (7.4)) Rifampicin: The concomitant administration of deferasirox tablets for oral suspension (single dose of 30 mg per kg) and the

rong uridine diphosphate glucuronosyltransferase (UGT) inducer rifampicin (600 mg per day for 9 days) decreased deferasirox AUC_{ist} by 44% (see Drug Interactions (7.5)). <u>Cholestyramine</u>: The concomitant administration of cholestyramine after a single dose of deferasirox tablets for oral suspension decreased deferasirox AUC., by 45% /see Drug Interactions (7.6). Busulfan: Concomitant administration of deferasirox and busulfan resulted in an increase of busulfan exposure (AUC).

In vitro Studies: Deferasirox inhibited human CYP2A6, CYP2D6, and CYP2C19 in vitro. Deferasirox is not a substrate of P-plycoprotein, MRP1 or MRP2. Pharmacokinetics in Specific Populations

deferasirox was less than in adult patients. In children less than 6 years of age, systemic exposure was about 50% lower than in adults. Sex: The apparent clearance is 17.5% lower in females compared to males. Renal Impairment: Compared to patients with MDS and eGFR greater than 60 mL/min/1.73 m², patients with MDS and eGFR 40 to 60 mL/min/1.73 m²(n = 34) had approximately 50% higher mean deferasirox trough plasma con

Hepatic Impairment: In a single dose (20 mg/kg) study in patients with varying degrees of hepatic impairment, deferasiron

irox increased 16% in 6 patients with mild (Child-Pugh A) hepatic impairment, and 76% in 6 patients with moderate (Child-Pugh B) hepatic impairment compared to 6 patients with normal hepatic function. The impact of severe (Child-Pugh C) rment was assessed in only 1 patient

13 NONCLINICAL TOXICOLOGY ${\bf 13.1} \quad \textbf{Carcinogenesis, Mutagenesis, Impairment of Fertility}$ A 104-week oral carcinogenicity study in Wistar rats showed no evidence of carcinogenicity from deferasirox at doses up to

60 mg/kg/day (0.7 times the MRHD on an mg/m² basis). A 26-week oral carcinogenicity study in p53 (+/-) transgenic mic ence of carcinogenicity from deferasirox at doses up to 200 mg/kg/day (1.2 times the MRHD on a mg/m² basis) in males and 300 mg/kg/day (1.7 times the MRHD on a mg/m²basis) in females Deferasirox was negative in the Ames test and chromosome aberration test with human peripheral blood lymphocytes. It

was positive in 1 of 3 in vivo oral rat micronucleus tests. Deferasirox at oral doses up to 75 mg/kg/day (0.9 times the MRHD on a mg/m²basis) was found to have no adverse effect on


sirox was evaluated in healthy subjects. There are no clinical data in patients with deferasirox. Deferasirox contains the same active ingredient as deferasirox tablets for oral suspension. The following information is based on clinical trials conducted with deferasirox tablets for oral suspension.

Transfusional Iron Overload The primary efficacy study, Study 1 (NCT00061750), was a multicenter, open-label, randomized, active-comparator control study to compare deferasirox tablets for oral suspension and deferoxamine in patients with beta-thalassemia and transfusional hemosiderosis. Patients greater than or equal to 2 years of age were randomized in a 1:1 ratio receive either oral deferasirox tablets for oral suspension at starting doses of 5, 10, 20, or 30 mg per kg once daily or subcutaneous deferoxamine at starting doses of 20 to 60 mg per kg for at least 5 days per week based on LIC at baseline (2 to 3, greater than 3 to 7, greater than 7 to 14, and greater than 14 mg Fe/g dry weight). Patients randomized to deferoxamine who had LIC values less than 7 mg Fe/g dry weight were permitted to continue on their prior deferoxamine dose, even though the dose

may have been higher than specified in the protocol. Patients were to have a liver biopsy at baseline and end of study (after 12 months) for LIC. The primary efficacy endpoint was defined as a reduction in LIC of greater than or equal to 3 mg Fe/g dry weight for baseline values greater than or equal to 10 mg Fe/g dry weight, reduction of baseline values between 7 and less than 10 to less than 7 mg Fe/g dry weight, or ance or reduction for baseline values less than 7 mg Fe/g dry weight. A total of 586 patients were randomized and treated, 296 with deferasirox tablets for oral suspension and 290 with

deferoxamine. The mean age was 17.1 years (range, 2 to 53 years); 52% were females and 88% were Caucasian. The primary efficacy population consisted of 553 patients (deferasirox tablets for oral suspension n=276; deferoxamine n=277) who had LIC evaluated at baseline and 12 months or discontinued due to an adverse reaction. The percentage of patients achieving the primary endpoint was 52.9% for deferasirox tablets for oral suspension and 66.4% for deferoxamine. The relative efficacy of deferasirox to deferoxamine cannot be determined from this study. In patients who had an LIC at baseline and at end of study, the mean change in LIC was $\cdot 2.4$ mg Fe/g dry weight in patients

 $treated \ with \ deferasirox\ tablets\ for\ oral\ suspension\ and\ -2.9\ mg\ Fe/g\ dry\ weight\ in\ patients\ treated\ with\ deferoxamine$ $Reduction of LIC and serum ferritin \,was observed \,with \,defer a sirox \,tablet \,for \,oral \,suspension \,doses \,of \,20\,to \,30\,mg \,per\,kg\,per\,kg \,per\,kg \,per\,k$ day. Deferasirox tablets for oral suspension doses below 20 mg per kg per day failed to provide consistent lowering of LII and serum ferritin levels (Figure 1). Therefore, a starting dose of 20 mg per kg per day is recommended [see Dosage and



Study 2 (NCT00061763) was an open-label, noncomparative trial of efficacy and safety of deferasirox tablets for oral suspension given for 1 year to patients with chronic anemias and transfusional hemosiderosis. Similar to Study 1, patient received 5, 10, 20, or 30 mg per kg per day of deferasirox tablets for oral suspension based on baseline LIC A total of 184 patients were treated in this study: 85 patients with beta-thalassemia and 99 patients with other congenital or acquired anemias (myelodysplastic syndromes, n – 47; Diamond-Blackfan syndrome, n – 30; other, n – 22). Nineteen percent (19%) of patients were less than 16 years of age and 16% were greater than or equal to 65 years of age. There was a reduction in the absolute LIC from baseline to end of study (-4.2 mg Fe/g dry weight).

Study 3 (NCT00067080) was a multicenter, open-label, randomized trial of the safety and efficacy of deferasirox tablets for oral suspension relative to deferoxamine given for 1 year in patients with sickle cell disease and transfusiona osiderosis. Patients were randomized to deferasirox tablets for oral suspension at doses of 5, 10, 20, or 30 mg per kg per day or subcutaneous deferoxamine at doses of 20 to 60 mg per kg per day for 5 days per week according to bas A total of 195 patients were treated in this study: 132 with deferasirox tablets for oral suspension and 63 with deferoxamine. Forty-four percent (44%) of patients were less than 16 years of age and 91% were black. At end of study, the mean change in LIC (as measured by magnetic susceptometry by a superconducting quantum interference device) in the per protocol-1 (PP-1) population, which consisted of patients who had at least 1 post-baseline LIC assessment, was -1.3 mg Fe/g dry weight for patients receiving deferasirox tablets for oral suspension (n = 113) and -0.7 mg Fe/g dry weight fo receiving deferoxamine (n = 54)

One-hundred five (105) patients with thalassemia major and cardiac iron overload were enrolled in a study assessing the change in cardiac magnetic resonance imaging (MRI) T2° value (measured in milliseconds, [ms]) before and after treatment with deferasirox. Cardiac T2° values at baseline ranged from 5 to less than 20 ms. The geometric mean of cardiac T2° in the 68 patients who completed 3 years of deferasirox tablets for oral suspension therapy increased from 11.98 ms at baseline to 17.12 ms at 3 years. Cardiac T2* values improved in patients with severe cardiac iron overload (less than 10 ms) and in those with mild to moderate cardiac iron overload (greater than or equal to 10 to less than 20 ms). The clinical significance o Six hundred twenty-seven (627) patients with MDS were enrolled across 5 uncontrolled trials. Two hundred thirty-nine of

the 627 patients were enrolled in trials that limited enrollment to patients with IPSS Low or Intermediate 1 risk MDS, and the remaining 388 patients were enrolled in trials that did not specify MDS risk stratification but required a life expectancy of greater than 1 year. Planned duration of treatment in these trials ranged from 1 year (365 patients) to 5 years (47 nts). These trials evaluated the effects of deferasirox tablets for oral suspension therapy on parameters overload, including LIC (125 patients) and serum ferritin (627 patients). The percent of patients completing planned duration of treatment was 51% in the largest 1-year study, 52% in the 3-year study and 22% in the 5-year study. The major causes for treatment discontinuation were withdrawal of consent, adverse reaction, and death. Over 1 year of follow-up across these pooled studies, mean change in serum ferritin was -332.8 (±2615.59) mcg/L (n – 593) and mean change in LIC was $-5.9 (\pm 8.32) \, mg \, Fe/g \, dw \, (n=68). \, Results \, of these pooled studies in 627 \, patients with MDS suggest a progressive decrease in serum ferritin and LIC beyond 1 year in those patients who are able to continue deferasirox tablets for oral suspension.$ Study 4 (TELESTO; NCT 00940602) was a randomized, double-blind, placebo-controlled trial performed in 225 patient with MDS (Low/Int-1 risk) and transfusional iron overload of which 149 were treated with deferasirox and 76 received

failure, liver function impairment, liver cirrhosis, or progression to acute myeloid leukemia; whichever occurred first Non-Transfusion-Dependent Thalassemia Study 5 (NCT00873041) was a randomized, double-blind, placebo-controlled trial of treatment with deferasirox tablets for oral suspension for patients 10 years of age or older with NTDT syndromes and iron overload. Eligible patients had an LIC of at least 5 mg Fe(g dw measured by RZ MRI and a serum ferritin exceeding 300 mcg/L at scz. English consecutive values at least 14 days apart from each other). A total of 166 patients were randomized, 55 to the deferasirox tablets for oral suspension 5 mg/kg/day dose group, 55 to the deferasirox tablets for oral suspension 10 mg/kg/day dose group, and 56 to placebo (28 to each matching placebo group). Doses could be increased after 6 months if the LIC receeded 7 mg Felg dw and the LIC reduction from baseline was less than 15%. The patients enrolled included 89 males and 77 females. The underlying disease was beta-thalassemia intermedia in 95 (57%) patients. HbE beta-thalassemia in 49 (30%) patients, and alpha halassemia in 22 (13%) patients. There were 17 pediatric patients in the study. Caucasians comprised 57% of the study population and Asians comprised 42%. The median baseline LIC (range) for all patients was 12.1 (2.6 to 49.1) mg Fe/g dw Follow-up was for 1 year. The primary efficacy endpoint of change in LIC from baseline to Week 52 was statistically significant in favor of both deferasirox dose groups compared with placebo (p less than or equal to 0.001) (Table 5).

Furthermore, a statistically significant dose effect of deferasirox was observed in favor of the 10 mg/kg/day dose group (10 versus 5 mg/kg/day, p = 0.009). In a descriptive analysis, the target LIC liess than 5 mg Fe/g dw) was reached by 15(27% of 55 patients in the 10 mg/kg/day arm, 8 (15%) of 55 patients in the 5 mg/kg/day arm and 2(4%) of 56 patients in the

Study 6 (NCT00873041) was an open-label trial of deferasirox tablets for oral suspension for the treatment of patients

placebo. The observed hazard ratio of 0.64 (95% Cl: 0.42, 0.96) suggests a positive impact of deferasirox on event-free survival (EFS, a composite endpoint defined as death, worsening cardiac function, hospitalization for congestive heart

previously enrolled on Study 5, including cross-over to active treatment for those previously treated with placebo. The starting dose of deferasirox tablets for oral suspension in Study 6 was assigned based on the patient's LIC at completion of

 $Study\ 5,\ being\ 20\ mg/kg/day\ for\ an\ LIC\ exceeding\ 15\ mg\ Fe/g\ dw,\ 10\ mg/kg/day\ for\ LIC\ 3\ to\ 15\ mg\ Fe/g\ dw,\ and\ observation$ if the LIC was less than 3 mg Fe/g dw. Patients could continue on 5 mg/kg/day if they had previously exhibited at least a 30% reduction in LIC. Doses could be increased to a maximum of 20 mg/kg/day after 6 months if the LIC was more than 7 mg Fe/g dw and the LIC reduction from baseline was less than 15%. The primary efficacy endpoint in Study 6 was the proportion of the LIC reduction from the LICpatients achieving an LIC less than 5 mg Fe/g dw. A total of 133 patients were enrolled. Twenty patients began Study 6 with an LIC less than 5 mg Fe/g dw. Of the 113 patients with a baseline LIC of at least 5 mg Fe/g dw in Study 6, the target LIC (less than 5 mg Fe/g dw) was reached by 39 patients (35%). The responders included 4 (10%) of 39 patients treated at 20 mg/kg/day for a baseline LIC exceeding 15 mg Fe/g dw, and 31 (51%) of 61 patients treated at 10 mg/kg/day for a baseline LIC between 5 and 15 mg Fe/g dw. The absolute change in LIC at Week 52 by starting dose is shown in Table 5 $^{\circ}$

Study 7 (NCT01709838) was an open-label, single-arm, multi-center, 5-year study to evaluate the efficacy and safety of deferasirox tablets for oral suspension in iron overloaded patients with NTDT of 10 years of age or older. All patients started treatment on 10 mg/kg/day deferasirox tablets for oral suspension for four weeks. At Week 4, dose escalation was based on baseline LIC. At Week 24 and every 6 months thereafter, further dose adjustments were made according to the LIC at that visit. Treatment was interrupted when LIC < 3 mg Fe/g dw or serum ferritin < 300 ng/mL and was re-started at 10 ng/kg/day when LIC ≥ 5 mg Fe/g dw and serum ferritin ≥ 300 ng/mL. Throughout the study, the maximum dose of deferasirox tablets for oral suspension given was 30 mg/kg/day.

A total of 134 patients were enrolled in the study. Eligible patients were required to have an LIC of at least 5 mg Fe/g dw measured by R2 MRI and a serum ferritin at least of 300 ng/mL at screening. The mean absolute change of LIC from E to Week 52 was -6.7 mg Fe/g dw. The reduction in LIC was sustained until Week 260 (5 years) with the mean absolute change in LIC from Baseline to Week 260 of \cdot 10.6 mg Fe/g dw. In the subset of patients with Baseline LIC > 15 mg Fe/g dw (49 patients), 51.0% achieved a first LIC < 5 mg Fe/g dw (95% CI: 37.5, 64.4) with a median time of 28.6 months. In the subset of patients with target LIC of < 3 mg Fe/g dw (61 patients), 39.3% developed first LIC \geq 5 mg Fe/g dw in the

Deferasirox Tablets for Oral Suspension Starting Dose ^a						
	Placebo	5 mg/kg/day	10 mg/kg/day	20 mg/kg/day		
Study 5 ^b						
Number of Patients	n = 54	n = 51	n = 54			
Mean LIC at Baseline (mg Fe/g dw)	16.1	13.4	14.4			
Mean Change (mg Fe/g dw)	+0.4	-2.0	-3.8			
(95% Confidence Interval)	(-0.6, +1.3)	(-2.9, -1.0)	(-4.8, -2.9)			
Study 6						
Number of Patients		n = 8	n = 77	n = 43		
Mean LIC at Baseline (mg Fe/g dw)		5.6	8.8	23.5		
Mean Change (mg Fe/g dw)		-1.5	-2.8	-9.1		
(95% Confidence Interval)		(-3.7, +0.7)	(-3.4, -2.2)	(-11.0, -7.3)		
Study 7						
Number of Patients			n = 127			
Mean LIC at Baseline (mg Fe/g dw)			15.1			
Mean Change (mg Fe/g dw)			-6.7			
(95% Confidence Interval)			(-7.9 -5.5)			

^b Least square mean change for Study 5.

Carton of 30 child-resistant sachets

Warnings and Precautions (5.3)].

Allergic Reactions

Randomized dose in Study 5 or assigned starting dose in Study 6 and Study 7.

HOW SUPPLIED/STORAGE AND HANDLING Deferasirox Oral Granules 90 mg are supplied as white to off white granular powder in sachet. They are available in Carton of 30 child-resistant sachets NDC 31722-029-32 Deferasirox Oral Granules 180 mg are supplied as white to off white granular powder in sachet. They are available in Carton of 30 child-resistant sachets NDC 31722-030-32

Deferasirox Oral Granules 360 mg are supplied as white to off white granular powder in sachet. They are available in

Store deferasirox oral granules at 20° C to 25° C (68° F to 77° F); excursions are permitted between 15° C to 30° C (59° F to 17 PATIENT COUNSELING INFORMATION Advise the patient to read the FDA-approved patient labeling (Medication Guide

Advise patients to take deferasirox oral granules by sprinkling the full dose on soft food (e.g., yogurt or applesauce)

NDC 31722-031-32

tial for kidney toxicity when taking deferasirox oral granules. Inform patients of the signs

immediately prior to use and administered orally. Advise patients to take deferasirox oral granules once a day, preferably at the same time each day. Deferasirox oral granules may be taken on an empty stomach or with a light meal (contains less than 7% fat content and approximately 250 calories). Examples of light meals include 1 whole wheat English muffin, 1 packet jelly (0.5 ounces), and skim milk (8 fluid ounces) or a turkey sandwich (2 oz. turkey on whole wheat bread w/lettuce, tomato

and 1 packet mustard). Advise patients that blood tests will be performed frequently to check for damage to kidneys, liver, or blood cells [see Warnings and Precautions (5.1, 5.2, 5.3, 5.4, 5.5)]. Acute Kidney Injury, Including Acute Renal Failure

and symptoms of kidney injury. Advise patients to contact their healthcare provider immediately if they experience any of these symptoms [see Warnings and Precautions (5.1)]. Henatic Toxicity and Failure on patients about the potential for hepatic toxicity when taking deferasirox oral granules. Inform patients of the signs

and symptoms of hepatic toxicity. Advise patients to contact their healthcare provider immediately if they experience any of

these symptoms (see Warnings and Precautions (5.2)). GI Ulceration and Hemorrhage ts about the potential for the development of GI ulcers or bleeding when taking deferasirox in combination with drugs that have ulcerogenic or hemorrhagic potential, such as NSAIDs, corticosteroids, oral bisphosphonates, or anticoagulants. Inform patients of the signs and symptoms of GI ulcers or bleeding. Advise patients to contact their healthcare provider for symptoms of heartburn but to seek immediate medical attention for symptoms of GI hemorrhage /see

immediate medical attention (see Warnings and Precautions (5.7)). Severe skin reactions have been reported in patients taking deferasirox oral granules. Inform patients of the signs and immediately and seek immediate medical attention [see Warnings and Precautions (5.8)].

Serious allergic reactions (which include swelling of the throat) have been reported in patients taking deferasirox, usually

Skin Rash Skin rashes may occur during deferasirox treatment. If the skin rash is severe, advise patients to stop taking deferasirox and Pediatric Patients with Acute Illness Instruct pediatric patients and their caregivers to contact their healthcare provider during episodes of acute illness,

especially if the patient has not been drinking fluids or the patient has volume depletion due to fever, vomiting, or diarrhea osure was increased compared to patients with normal hepatic function. The average total (free and bound) AUC., of Auditory and Ocular Testing cause auditory and ocular disturbances have been reported with deferasirox, conduct auditory testing and ophthalmic testing before starting deferasirox treatment and thereafter at regular intervals. Advise patients to contact their healthcare

> provider if they develop visual or auditory changes during treatment/see Warnings and Precautions (5.10)]. **Drug Interactions** Caution patients not to take aluminum containing antacids and deferasirox oral granules simultaneously [see Drug Caution patients about potential loss of effectiveness of drugs metabolized by CYP3A4 (e.g., cyclosporine, simvastatir $hormonal\ contraceptive\ agents)\ when\ deferasirox\ is\ administered\ with\ these\ drugs\ \textit{/see}\ \textit{Drug}\ \textit{Interactions}\ \textit{(7.2)/}.$

inducers (e.g., rifampicin, phenytoin, phenobarbital, ritonavir). Based on serum ferritin levels and clinical respons

increases in the dose of deferasirox when concomitantly used with potent UGT inducers [see Drug Interactions (7.5)]. Caution patients about potential loss of effectiveness of deferasirox when administered with drugs that are bile acid

sequestrants (e.g., cholestyramine, colesevelam, colestipol). Based on serum ferritin levels and clinical response, consider ncreases in the dose of deferasirox when concomitantly used with bile acid sequestrants [see Drug Interactions (7.6)]. Caution patients with diabetes to monitor their glucose levels more frequently when repaglinide is used concomitantly with

Advise patients to store deferasirox oral granules in a dry, room-temperature environment (see How Supplied/Storage and

Caution nations about notential loss of effectiveness of deferasirox when administered with drugs that are notent UGT

Driving and Using Machines Caution patients experiencing dizziness to avoid driving or operating machinery [see Adverse Reactions (6.1)]



Piscataway, NJ 08854

Handling Instructions

By: Annora Pharma Pyt. Ltd.

2102760